

Polytopes, Hopf algebras and Quasi-symmetric functions

Victor M. Buchstaber

Nickolai Erokhovets

Abstract

In this paper we use the technique of Hopf algebras and quasi-symmetric functions to study the combinatorial polytopes. Consider the free abelian group $\mathcal{P}$ generated by all combinatorial polytopes. There are two natural bilinear operations on this group defined by a direct product $\times$ and a join $\ast$ of polytopes. It turns out that $(\mathcal{P}, \times)$ is a commutative associative bigraded ring of polynomials with graduations $(2n, 2(m-n))$, where n is the dimension, and m is the number of facets of a polytope. $(\mathcal{P}, \ast)$ is a commutative associative ring without a unit. If we add the empty set $\emptyset$ as a (-1) -dimensional polytope, then it gives a unit, and we obtain the commutative associative threegraded ring of polynomials $\mathcal{RP}$ with graduations $(2(n+1), 2(v-n-1), 2(m-n-1))$, where v is the number of vertices of a polytope. The ring $\mathcal{RP}$ has the structure of a graded Hopf algebra and the natural correspondence $P \rightarrow L(P)$ that sends a polytope to its face lattice embeds $\mathcal{RP}$ into the Rota-Hopf algebra $\mathcal{R}$ of graded posets as a Hopf subalgebra. It turns out that $\mathcal{P}$ has a natural Hopf comodule structure over the Hopf algebra $\mathcal{RP}$. Faces operators d_k that send a polytope to the sum of all its $(n-k)$ -dimensional faces define on both rings the Hopf module structures over the Hopf algebra $\mathcal{U}$ that is universal in the category of Leibnitz-Hopf algebras with the antipode $U_k \rightarrow (-1)^k U_k$. This structure gives a ring homomorphism $R \rightarrow Qsym \otimes R$, where R is $\mathcal{P}$ or $\mathcal{RP}$. Composing this homomorphism with the characters $P^n \rightarrow \alpha^n$ of $\mathcal{P}$, $P^n \rightarrow \alpha^{n+1}$ of $\mathcal{RP}$, and with the counit we obtain the ring homomorphisms $f: \mathcal{P} \rightarrow Qsym[\alpha]$, $f_{\mathcal{RP}}: \mathcal{RP} \rightarrow Qsym[\alpha]$, and $F^*: \mathcal{RP} \rightarrow Qsym$, where F is the Ehrenborg transformation. We describe the images of these homomorphisms in terms of functional equations, prove that these images are rings of polynomials over $\mathbb{Q}$, and find the relations between the images, the homomorphisms and the Hopf comodule structures. For each homomorphism f , $f_{\mathcal{RP}}$, and F the images of two polytopes coincide if and only if they have equal flag f -vectors. Therefore algebraic structures on the images give the information about flag f -vectors of polytopes. The homomorphism f is an isomorphism on the graded group BB generated by the polytopes introduced by M. Bayer and L. Billera to find the linear span of flag f -vectors of convex polytopes. This gives the group BB a structure of the ring isomorphic to $f(\mathcal{P})$. Developing this approach one can build new combinatorial invariants of convex polytopes that can not be expressed in terms of flag numbers.

1 Introduction

Convex polytopes is a classical object of convex geometry. In recent times the solution of problems of convex geometry involves results of algebraic geometry and topology, commutative and homological algebra. There are remarkable results lying on the crossroads of the polytope theory, theory of complex manifolds, equivariant topology and singularity theory (see the survey [BR]). The example of successful collaboration of the polytope theory and differential equations can be found in [Buch].

To study the combinatorics of convex polytopes we consider the free abelian group $\mathcal{P}$ generated by all combinatorial convex polytopes. It turns out that this object has a rich algebraic nature. The direct product $\times$ defines on $\mathcal{P}$ the structure of a bigraded associative commutative ring $(\mathcal{P}, \times)$ with graduations $(2n, 2(m-n))$, where n is the dimension, and m is the number of facets of a polytope. The bigraded structure gives the decomposition of $\mathcal{P}$ into a direct sum of finitely generated free abelian groups

$$\mathcal{P} = \mathcal{P}^0 + \sum_{m \geq 2} \sum_{n=1}^{m-1} \mathcal{P}^{2n, 2(m-n)}.$$

The duality operator $*$ defines on $\mathcal{P}$ another multiplication $P \circ Q = (P^* \times Q^*)^*$.

The join of two polytopes $P \ast Q$ is a bilinear operation of degree $+2$. Then $(\mathcal{P}, \ast)$ is a commutative associative ring without a unit. Set $\mathcal{RP} = \mathbb{Z}\emptyset \oplus \mathcal{P}$. The empty set $\emptyset$ may be considered as a (-1) -

dimensional polytope such that $\emptyset * P = P$. Then the free abelian group $\mathcal{RP}$ has the structure of a threegraded commutative associative ring with graduations $(2(n+1), 2(v-n-1), 2(m-n-1))$, where v is the number of vertices of a polytope. It turns out that $\mathcal{RP}$ is a graded Hopf algebra with graduation $2(n+1)$ and the comultiplication

$$\Delta(P) = \sum_{\emptyset \subseteq F \subseteq P} F \otimes (P/F)$$

The correspondence $P \rightarrow L(P)$, where $L(P)$ is a face lattice of a polytope P defines an embedding of $\mathcal{RP}$ as a Hopf subalgebra into the Hopf algebra $\mathcal{R}$ of graded posets introduced by Joni and Rota in [JR].

In the focus of our interest we put the ring of linear operators $\mathcal{L}(\mathbf{R}) = \text{Hom}_{\mathbb{Z}}(\mathbf{R}, \mathbf{R})$, where $\mathbf{R} = \mathcal{P}$ or $\mathcal{RP}$, and its subring $\mathcal{D}$ generated by the operators d_k , $k \geq 1$ that map an n -dimensional polytope to the sum of all its $(n-k)$ -dimensional faces. $d = d_1$ is a derivation, so $\mathbf{R}$ is a differential ring.

The ring $\mathcal{D}$ has a canonical structure of a Leibnitz-Hopf algebra (see Section 3.2). We prove that $\mathcal{D}$ is isomorphic to the universal Hopf algebra in the category of Leibnitz-Hopf algebras with the antipode $\chi(d_k) = (-1)^k d_k$.

$$\mathcal{D} \simeq \mathcal{Z}/J_{\mathcal{D}}$$

where $\mathcal{Z} = \mathbb{Z}\langle Z_1, Z_2, \dots \rangle$ is the universal Leibnitz-Hopf algebra – the free associative Hopf algebra with the comultiplication

$$\Delta Z_k = \sum_{i+j=k} Z_i \otimes Z_j, \quad Z_0 = 1,$$

and $J_{\mathcal{D}}$ is the two-sided Hopf ideal generated by the relations $\sum_{i+j=n} (-1)^i Z_i Z_j = 0$, $n \geq 1$.

This algebra appears in various application of theory of Hopf algebras in combinatorics: over the rationals it is isomorphic to the graded dual of the odd subalgebra $S_-(\text{Qsym}[t_1, t_2, \dots], \zeta_Q)$ [ABS, Remark 6.7], to the algebra of forms on chain operators [BLiu, Proposition 3.2], to the factor algebra of the algebra of Piery operators on a Eulerian poset by the ideal generated by the Euler relations [BMSW, Example 5.3].

The action of $\mathcal{D}$ on $\mathbf{R}$ satisfies the property $D_{\omega}(PQ) = \mu \circ (\Delta D_{\omega})(P \otimes Q)$, so the ring $\mathbf{R}$ has the structure of a Milnor module over $\mathcal{D}$.

The graded dual Hopf algebra to $\mathcal{Z}$ is the algebra of quasi-symmetric functions $\text{Qsym}[t_1, t_2, \dots]$. The character $\xi_{\alpha}: \mathcal{P} \rightarrow \mathbb{Z}[\alpha]$, $P^n \rightarrow \alpha^n$ induces the homomorphism $\mathcal{P} \rightarrow \mathcal{D}^*[\alpha]$, and the composition with the embedding $\mathcal{D}^* \subset \text{Qsym}[t_1, t_2, \dots]$ gives the ring homomorphism

$$f: \mathcal{P} \rightarrow \text{Qsym}[t_1, t_2, \dots][\alpha]$$

We call $f(\alpha, t_1, t_2, \dots)$ the generalized f -polynomial. It can be shown that $f(\alpha, t_1, 0, 0, \dots) = f_1(\alpha, t)$ is a homogeneous f -polynomial in two variables introduced in [Buch]. For simple polytopes we have $f(\alpha, t_1, t_2, \dots) = f_1(\alpha, t_1 + t_2 + \dots)$.

The homomorphism f can be considered in a more general context. The structure of a right Milnor module on the ring $\mathbf{R}$ over the universal Hopf algebra $\mathcal{Z}$ defines a left Hopf comodule structure $\mathbf{R} \rightarrow \text{Qsym} \otimes \mathbf{R}$. Then any ring homomorphism $\mathbf{R} \rightarrow \mathcal{R}_1$ defines the ring homomorphism $\mathbf{R} \rightarrow \text{Qsym} \otimes \mathcal{R}_1$. In particular, the character $P^n \rightarrow \alpha^n$ defines the ring homomorphism $f: \mathcal{P} \rightarrow \text{Qsym}[\alpha]$, the character $\varepsilon_{\alpha}(P^n) = \alpha^{n+1}$ defines a ring homomorphism $f_{\mathcal{RP}}: \mathcal{RP} \rightarrow \text{Qsym}[\alpha]$, the counit ε defines a ring homomorphism $F^*: \mathcal{RP} \rightarrow \text{Qsym}$ such that $F(P)$ is the Ehrenborg F -quasi-symmetric function of a graded poset $L(P)$. The image of two polytopes under any of homomorphisms $f, f_{\mathcal{RP}}$, or F^* coincide if and only if the polytopes have equal flag f -numbers.

It turns out that there is a nice algebraic structure including the rings $\mathcal{P}$ and $\mathcal{RP}$. Namely, the formula

$$\Delta_{\mathcal{RP}} P = \sum_{F \subseteq P} F \otimes (P/F)$$

defines on the ring $\mathcal{P}$ the structure of a graded Hopf comodule over the Hopf algebra $\mathcal{RP}$ with respect to graduations defined by dimension of a polytope. This approach clarifies the structures mentioned above and allows us to build new ring homomorphisms from the ring of polytopes $\mathcal{P}$ to rings of polynomials such that the image of a polytope is not determined by its flag f -vector. On this way one can build new

combinatorial invariants of polytopes that can not be expressed in terms of flag numbers. This will be the topic of our next article.

The paper has the following structure:

Section 2 describes the main definitions and constructions involving convex polytopes.

- part 2.1 contains the main definitions and considers the basic constructions involving polytopes;
- in part 2.2 the ring of polytopes $\mathcal{P}$ is defined.
- in part 2.3 we establish the structure of a graded Hopf algebra on the ring $\mathcal{RP}$ and consider it as a Hopf subalgebra of the Rota-Hopf algebra $\mathcal{R}$.
- part 2.4 is devoted to linear operators defined on the rings $\mathcal{P}$ and $\mathcal{RP}$. In particular, the faces operators d_k are defined.
- part 2.5 recalls important facts about flag f -vectors and emphasizes some peculiarities we need in this paper.

Section 3 is devoted to necessary definitions and important facts about quasi-symmetric functions and Hopf algebras.

- part 3.1 contains the definition of quasi-symmetric functions;
- part 3.2 includes the definition of Leibnitz-Hopf algebras;
- part 3.3 – of Lie-Hopf algebras;
- part 3.4 describes the idea of the proof of the shuffle-algebra theorem.

Section 4 contains topological realizations of the Hopf algebras we study.

In Section 5 the structure theorem for the ring $\mathcal{D}$ is proved.

In Section 6 we define and study the generalized f -polynomial.

- part 6.1 contains the construction of f ;
- in part 6.2 we find the functional equations that describe the image of the ring $\mathcal{P}$ in $\text{Qsym}[t_1, t_2, \dots][\alpha]$. These equations are equivalent to the generalized Dehn-Sommerville equations discovered by M. Bayer and L. Billera in [BB].
- part 6.3 describes the properties that define the generalized f -polynomial in a unique way.

In Section 7 we study the Hopf algebra $\mathcal{D}^*$ as a Hopf subalgebra in Qsym .

- in part 7.1 the main theorem is proved;
- part 7.2 contains the applications of the main theorem. In particular, the image of the ring of polytopes $\mathcal{P}$ in $\mathcal{D}^*[\alpha]$ under the natural mapping φ_α is found.

In Section 8 the multiplicative structure of the ring $f(\mathcal{P} \otimes \mathbb{Q})$ is described. We prove that $f(\mathcal{P} \otimes \mathbb{Q})$ is a free polynomial algebra with dimension of the $2n$ -th graded component equal to the n -th Fibonacci number c_n ($c_0 = c_1 = 1$, $c_{n+1} = c_n + c_{n-1}$, $n \geq 1$). This gives the decomposition of the Fibonacci series into the infinite product

$$\frac{1}{1-t-t^2} = \sum_{n=0}^{\infty} c_n t^n = \prod_{i=1}^{\infty} \frac{1}{(1-t^i)^{k_i}},$$

where k_i is the number of generators of degree $2i$. The infinite product converges absolutely in the interval $|t| < \frac{\sqrt{5}-1}{2}$. The numbers k_n satisfy the inequalities $k_{n+1} \geq k_n \geq N_n - 2$, where N_n is the number of decompositions of n into the sum of odd numbers. Moreover, standard methods allow one to find the explicit formula for k_n , which in the case of prime p has the form

$$k_p = \sum_{j=1}^{\lfloor \frac{p}{2} \rfloor} \frac{\binom{p-j}{j}}{p-j}$$

In Section 9 we introduce the multiplicative structure on the graded group BB generated by the Bayer-Billera polytopes arising from the isomorphism with the ring $f(\mathcal{P})$.

Nowadays Hopf algebras is one of the central tools in combinatorics. There is a well-known Hopf algebra $\mathcal{R}$ of posets introduced by Joni and Rota in [JR]. Various aspects of this algebra were studied in [Ehr, ABS, Sch1, Sch2]. The generalization of the Rota-Hopf algebra was proposed in [RS]. In [Ehr] R. Ehrenborg introduced the F -quasi-symmetric function, which gives a Hopf algebra homomorphism from the Rota-Hopf algebra to $\text{Qsym}[t_1, t_2, \dots]$. Section 10 contains the following results:

- in part 10.1 we consider the Hopf comodule structures arising from the Hopf module structures on the rings $\mathcal{P}$ and $\mathcal{RP}$ over the Hopf algebras $\mathcal{Z}$ and $\mathcal{D}$;
- in part 10.2 we describe the ring homomorphisms that arise from this structures. As an example, we show how the generalized f -polynomial and the Ehrenborg F -quasi-symmetric function appear in this context;
- part 10.3 contains the definition and the main properties of the natural Hopf comodule structure on the ring $\mathcal{P}$ over the Hopf algebra $\mathcal{RP} \subset \mathcal{R}$;
- part 10.4 describes the interrelations between the Hopf comodule structures and the homomorphisms arising in this section. In particular, it is shown that the Hopf comodule structures from parts 10.1 and 10.3 are related by the Ehrenborg transformation.

Section 11 is devoted to the cone and the bipyramid operators C and B used by M. Bayer and L. Billera [BB] to find the linear span of flag f -vectors of polytopes. We show that the operations C and B can be defined on the ring $\text{Qsym}[t_1, t_2, \dots][\alpha]$ in such a way that $f(CP) = Cf(P)$ and $f(BP) = Bf(P)$. The same idea works for the ring $\mathcal{RP}$ and the homomorphisms $f_{\mathcal{RP}}$, and F^* .

- in part 11.1 we consider the case of $\mathcal{P}$;
- in part 11.2 – the case of $\mathcal{RP}$.

Section 12 deals with the problem of the description of flag f -vectors of polytopes. For simple polytope g -theorem gives the full description of the set of flag f -vectors. As it is mentioned in [Z1] even for 4-dimensional non-simple polytopes the corresponding problem is extremely hard. We show how this problem can be stated in terms of the ring of polytopes.

This article contains the results and corrections of some inaccuracies made in the work [BuchE].

2 Polytopes

2.1 Definitions and constructions

This section contains main definitions and constructions from the polytope theory. For details see the books [Gb], [Z2], [BP].

There are two algorithmically different ways to define a *convex polytope*.

Definition 2.1. A $\mathcal{V}$ -polytope is a convex hull of a finite set of points in some $\mathbb{R}^d$.

Definition 2.2. An $\mathcal{H}$ -polyhedron is an intersection of finitely many closed halfspaces in some $\mathbb{R}^d$. An $\mathcal{H}$ -polytope is a bounded $\mathcal{H}$ -polyhedron.

The classical fact that these two definitions are equivalent is proved in different books, for example in [Z2, Theorem 1.1].

The *dimension* of a convex polytope is the dimension of its affine hull. Without loss of generality in most cases we will assume that an n -dimensional polytope P lies in the space $\mathbb{R}^n$.

It is not difficult to prove the following lemma:

Proposition 2.3. Any n -dimensional convex polytope P^n up to a translation can be represented in the form:

$$P^n = \{\mathbf{x} \in \mathbb{R}^n : \langle \mathbf{a}_i, \mathbf{x} \rangle + 1 \geq 0, i = 1, \dots, m\}, \quad (1)$$

where the system of inequalities is irredundant, that is the absence of any one of them changes the set P^n .

In this case the polytope P^n has exactly m facets

$$F_i = P^n \cap \{\mathbf{x} \in \mathbb{R}^n : \langle \mathbf{a}_i, \mathbf{x} \rangle + 1 = 0\}, \quad i = 1, \dots, m.$$

Proof. It is enough to make by a translation the origin $\mathbf{0}$ the inner point of P . $\square$

Two polytopes P^n and Q^n are said to be *combinatorially equivalent* if there exists a bijection between their face lattices $L(P)$ and $L(Q)$ that preserves an inclusion relation. A *combinatorial polytope* is an equivalence class of combinatorially equivalent convex polytopes.

The notion of the general position from the point of view of two different definitions of a convex polytope give two special classes of polytopes.

Definition 2.4. A polytope P^n is called *simple* if any vertex of P belongs to exactly n facets.

A polytope P^n is called *simplicial* if any facet of P is an $(n - 1)$ -dimensional simplex (i.e. contains exactly n vertices).

Definition 2.5 (dual polytope). Let P^n be an n -dimensional polytope (1). Then a *dual* (or *polar*) polytope is defined as

$$(P^n)^* = \{\mathbf{y} \in \mathbb{R}^n : \langle \mathbf{y}, \mathbf{x} \rangle + 1 \geq 0, \forall \mathbf{x} \in P^n\}.$$

It can be shown (see, e.g. [Z2]) that

$$(P^n)^* = \text{conv}\{\mathbf{a}_i, i = 1, \dots, m\},$$

and $(P^*)^* = P$.

There is a bijection $F \longleftrightarrow F^\diamond$ between the i -faces of P and the $(n - 1 - i)$ -faces of P^* such that

$$F \subset G \Leftrightarrow G^\diamond \subset F^\diamond.$$

For any simple polytope P its polar polytope P^* is simplicial and vice versa.

Definition 2.6 (face polytope). For the face F of the polytope P let us define a *face polytope* P/F as

$$P/F = (F^\diamond)^*$$

It is easy to see that the face polytope has dimension $\dim P - \dim F - 1$ and the face lattice

$$L(P/F) = [F, P] = \{G \in L(P) : F \subseteq G \subseteq P\}.$$

Example 2.7. Let P^n be a simple polytope. Then for any proper k -dimensional face F of P we have $P/F = \Delta^{n-k-1}$.

Definition 2.8 (direct product). For two polytopes $P^{n_1} \subset \mathbb{R}^{n_1}$ and $Q^{n_2} \subset \mathbb{R}^{n_2}$ with m_1 and m_2 facets respectively the *direct product*

$$P^{n_1} \times Q^{n_2} = \{(\mathbf{x}, \mathbf{y}) \in \mathbb{R}^{n_1} \times \mathbb{R}^{n_2} : \mathbf{x} \in P^{n_1}, \mathbf{y} \in Q^{n_2}\}$$

is an $(n_1 + n_2)$ -dimensional convex polytope. Its faces are direct products of faces of P^{n_1} and Q^{n_2} . In particular, it has $m_1 + m_2$ facets.

Definition 2.9 (operation $\circ$). For two polytopes

$$\begin{aligned} P^{n_1} &= \{\mathbf{x} \in \mathbb{R}^{n_1} : \langle \mathbf{a}_i, \mathbf{x} \rangle + 1 \geq 0, i = 1, \dots, m_1\}, \\ Q^{n_2} &= \{\mathbf{y} \in \mathbb{R}^{n_2} : \langle \mathbf{b}_j, \mathbf{y} \rangle + 1 \geq 0, j = 1, \dots, m_2\} \end{aligned}$$

let us define the polytope

$$P^{n_1} \circ Q^{n_2} = \text{conv}(P^{n_1} \times \{\mathbf{0}\} \cup \{\mathbf{0}\} \times Q^{n_2}) \subset \mathbb{R}^{n_1} \times \mathbb{R}^{n_2}$$

Proposition 2.10. For two polytopes P^{n_1} and Q^{n_2} represented in the form (1) we have

$$(P^{n_1} \times Q^{n_2})^* = (P^{n_1})^* \circ (Q^{n_2})^*$$

Proof. The polytope $P^{n_1} \times Q^{n_2}$ is defined by the inequalities

$$\langle \mathbf{a}_i, \mathbf{x} \rangle + 1 \geq 0, i = 1, \dots, m_1, \quad \langle \mathbf{b}_j, \mathbf{y} \rangle + 1 \geq 0, j = 1, \dots, m_2,$$

therefore,

$$(P^{n_1} \times Q^{n_2})^* = \text{conv}\{(\mathbf{a}_i, \mathbf{0}), (\mathbf{0}, \mathbf{b}_j), i = 1, \dots, m_1, j = 1, \dots, m_2\} = (P^{n_1})^* \circ (Q^{n_2})^*$$

□

Since the product of two simple polytopes is again a simple polytope, we see that for any two simplicial polytopes P and Q the polytope $P \circ Q$ is again simplicial.

Example 2.11 (bipyramid). Let us define a *bipyramid* as $BP = I \circ P$. Then we have $(BP)^* = I \times P^*$.

Definition 2.12 (join). For the space $\mathbb{R}^n$ let Δ^{n-1} be the regular simplex $\text{conv}(e_1, \dots, e_n)$, where $e_i = (0, \dots, 0, 1, 0, \dots, 0)$ is the i -th basis vector. Let $P^{n_1} \subset \mathbb{R}^{n_1+1}$, and $Q^{n_2} \subset \mathbb{R}^{n_2+1}$ be polytopes lying in the regular simplices Δ^{n_1} and Δ^{n_2} . A *join* $P * Q$ is defined as

$$P * Q = \text{conv}(P \times \{\mathbf{0}\} \cup \{\mathbf{0}\} \times Q) \subset \mathbb{R}^{n_1+1} \times \mathbb{R}^{n_2+1}$$

It is easy to see that the polytope $P * Q$ lies in the regular simplex $\Delta^{n_1+n_2+1}$.

From this construction is evident that the join is an associative operation.

Remark 2.13. In the definition of the join we used the polytopes that lie in the regular simplices. In fact, the affine type of $P * Q$ does not depend on the positions of P and Q in the regular simplices, moreover, it depends only on the affine types of P and Q .

Indeed, let $\{\mathbf{v}_1, \dots, \mathbf{v}_{n_1+1}\}$ be the set of vertices of P in general position, and $\{\mathbf{w}_1, \dots, \mathbf{w}_{n_2+1}\}$ be the set of vertices of Q in general position. Since the origin $O = \mathbf{0} \in \mathbb{R}^{n_1+1} \times \mathbb{R}^{n_2+1}$ does not belong to $\text{aff } P$, the vectors $\overrightarrow{O\mathbf{v}_1}, \dots, \overrightarrow{O\mathbf{v}_{n_1+1}}$ are linearly independent. Similarly the vectors $\overrightarrow{O\mathbf{w}_1}, \dots, \overrightarrow{O\mathbf{w}_{n_2+1}}$ form a basis in $\mathbb{R}^{n_2+1}$. Then the vectors $\overrightarrow{O\mathbf{v}_1}, \dots, \overrightarrow{O\mathbf{v}_{n_1+1}}, \overrightarrow{O\mathbf{w}_1}, \dots, \overrightarrow{O\mathbf{w}_{n_2+1}}$ are linearly independent in $\mathbb{R}^{n_1+1} \times \mathbb{R}^{n_2+1}$, and the set of points $\mathbf{v}_1, \dots, \mathbf{v}_{n_1+1}, \mathbf{w}_1, \dots, \mathbf{w}_{n_2+1}$ is in general position. This implies that $\dim P^{n_1} * Q^{n_2} = n_1 + n_2 + 1$. Now for any two realizations $(P \in \Delta^{n_1}, Q \in \Delta^{n_2})$ and $(P' \in \Delta^{n_1}, Q' \in \Delta^{n_2})$ the correspondence $\mathbf{v}_i \rightarrow \mathbf{v}'_i, \mathbf{w}_j \rightarrow \mathbf{w}'_j$ defines an affine isomorphism $\text{aff } P * Q \simeq \text{aff } P' * Q'$ such that $P \rightarrow P', Q \rightarrow Q'$. Then $\text{conv}(P \cup Q) \rightarrow \text{conv}(P' \cup Q')$, so $P * Q \simeq P' * Q'$. Since the affine types of P and Q are uniquely defined by the sets of points $\{\mathbf{v}_i\}$ and $\{\mathbf{w}_j\}$, we see, that the affine type of $P * Q$ depends only on the affine types of P and Q .

Proposition 2.14. $P^{n_1} * Q^{n_2}$ is an $(n_1 + n_2 + 1)$ -dimensional polytope.

Faces of $P * Q$ up to an affine equivalence are exactly $F * G$, where $\emptyset \subseteq F \subseteq P, \emptyset \subseteq G \subseteq Q$ are faces of P and Q respectively.

Proof. For the proof see Appendix A. □

Corollary 2.15. The face lattice $L(P * Q)$ is the direct product of the face lattices $L(P)$ and $L(Q)$. Therefore the join operation is well defined on combinatorial polytopes.

Remark 2.16. The join $P * Q$ can be defined in more invariant terms – as the convex hull of P and Q provided they are placed into affine subspaces of some $\mathbb{R}^N$ such that their affine hulls $\text{aff}(P)$ and $\text{aff}(Q)$ are skew.

Example 2.17. For the simplices Δ^k and Δ^l we have $\Delta^k * \Delta^l = \Delta^{k+l+1}$.

Proposition 2.18. For two combinatorial polytopes P and Q

$$(P * Q)^* = P^* * Q^*$$

Proof. Indeed, the faces of the polytope $(P * Q)^*$ have the form $(F * G)^\diamond$, and $(F * G)^\diamond \subseteq (F' * G')^\diamond$ if and only if $F' * G' \subseteq F * G$, i.e. $F' \subseteq F, G' \subseteq G$, which is equivalent to $F^\diamond \subseteq (F')^\diamond, G^\diamond \subseteq (G')^\diamond$. Therefore the correspondence $(F * G)^\diamond \longleftrightarrow F^\diamond * G^\diamond$ is a combinatorial equivalence of the polytopes $(P * Q)^*$ and $P^* * Q^*$. □

Example 2.19 (cone). Let us define a *cone* CP as $\text{pt} * P$, where pt is a point. Then $(CP)^* = C(P^*)$.

2.2 Ring of polytopes

Definition 2.20. Denote by $\mathcal{P}^{2n}$ the free abelian group generated by all n -dimensional combinatorial polytopes. For $n > 1$ we have the direct sum

$$\mathcal{P}^{2n} = \sum_{m \geq n+1} \mathcal{P}^{2n, 2(m-n)},$$

where $P^n \in \mathcal{P}^{2n, 2(m-n)}$ if it is a polytope with m facets, and $\text{rank } \mathcal{P}^{2n, 2(m-n)} < \infty$ for any fixed n and m such that $n < m$. The product of polytopes $P \times Q$ turns the direct sum

$$\mathcal{P} = \sum_{n \geq 0} \mathcal{P}^{2n} = \mathcal{P}^0 + \sum_{m \geq 2} \sum_{n=1}^{m-1} \mathcal{P}^{2n, 2(m-n)}$$

into a bigraded commutative associative ring, a *ring of polytopes*. The unit is $P^0 = \text{pt}$, a point.

Let us denote $\mathcal{P}^{[2n]} = \mathcal{P}^0 \oplus \mathcal{P}^2 \oplus \dots \oplus \mathcal{P}^{2n}$.

The direct product $P \times Q$ of simple polytopes P and Q is a simple polytope as well. Thus the abelian subgroup $\mathcal{P}_s \subset \mathcal{P}$ generated by all simple polytopes is a subring in $\mathcal{P}$.

Definition 2.21. A polytope P^n is called *indecomposable* if it can not be decomposed into a direct product $P_1 \times P_2$ of two polytopes of positive dimensions.

Proposition 2.22. $\mathcal{P}$ is a polynomial ring generated by indecomposable combinatorial polytopes.

For the proof see Appendix B.

Since the polytope $P = P_1 \times P_2$ is simple if and only if the polytopes P_1 and P_2 are simple, $\mathcal{P}_s$ is a polynomial ring generated by all indecomposable simple polytopes.

Since $(P^*)^* = P$, the correspondence $P \rightarrow P^*$ defines an involutory linear isomorphism of the ring $\mathcal{P}$.

Any isomorphism of graded abelian groups $A: \mathcal{P} \rightarrow \mathcal{P}$ defines a new multiplication

$$P \times_A Q = A^{-1}(AP \times AQ)$$

such that A is a graded ring isomorphism $(\mathcal{P}, \times_A) \rightarrow (\mathcal{P}, \times)$.

In particular, $*$ defines the multiplication

$$P \circ Q = (P^* \times Q^*)^*$$

Definition 2.23. Let us define an *alpha-character* $\xi_\alpha: \mathcal{P} \rightarrow \mathbb{Z}[\alpha]$:

$$\xi_\alpha(P^n) = \alpha^n$$

2.3 The Rota-Hopf algebra

Definition 2.24. Let P be a finite poset with a minimal element $\hat{0}$ and a maximal element $\hat{1}$.

An element y in P *covers* another element x in P if $x < y$ and there is no z in P such that $x < z < y$.

A poset P is called *graded* if there exists a rank function $\rho: P \rightarrow \mathbb{Z}$ such that $\rho(\hat{0}) = 0$ and $\rho(y) = \rho(x) + 1$ if y covers x . It is easy to see that if a rank function exists, then it is determined in a unique way.

Set $\rho(P) = \rho(\hat{1})$, and $\deg(P) = 2\rho(P)$.

Two finite graded posets are isomorphic if there exists an order preserving bijection between them.

Let us denote by $\mathcal{R}$ the graded free abelian group with basis the set of all isomorphism classes of finite graded posets.

$\mathcal{R}$ has the structure of a *graded connected Hopf algebra*:

- The multiplication $P \cdot Q$ is a cartesian product $P \times Q$ of posets P and Q : let $x, u \in P$, and $y, v \in Q$. Then

$$(x, y) \leq (u, v) \text{ if and only if } x \leq u \text{ and } y \leq v$$

- The unit element in $\mathcal{R}$ is the poset with one element $\hat{0} = \hat{1}$
- The comultiplication is

$$\Delta(P) = \sum_{\hat{0} \leq z \leq \hat{1}} [\hat{0}, z] \otimes [z, \hat{1}],$$

where $[x, y]$ is the subposet $\{z \in P \mid x \leq z \leq y\}$.

- The counit ε is

$$\varepsilon(P) = \begin{cases} 1, & \text{if } \hat{0} = \hat{1}; \\ 0, & \text{else} \end{cases}$$

- The antipode χ is

$$\chi(P) = \sum_{k \geq 0} \sum_{C_k} (-1)^k [x_0, x_1] \cdot [x_1, x_2] \cdots [x_{k-1}, x_k],$$

where $C_k = (\hat{0} = x_0 < x_1 < \cdots < x_k = \hat{1})$

The Hopf algebra of graded posets was originated in the work by Joni and Rota [JR]. Variations of this construction were studied in [Ehr, ABS, Sch1, Sch2]. The generalization of this algebra can be found in [RS].

Example 2.25. The simplest Boolean algebra $B_1 = \{\hat{0}, \hat{1}\}$ is the face lattice of the point pt.

$$\begin{aligned} \Delta(B_1) &= 1 \otimes B_1 + B_1 \otimes 1 \\ \chi(B_1) &= -B_1 \end{aligned}$$

Definition 2.26. There is a natural involutory graded ring isomorphism $*$ of the ring $\mathcal{R}$. For a graded poset P let P^* be a graded poset consisting of the same elements but with the reverse inclusion relation, i.e. $x \leq_P y$ if and only is $x \geq_{P^*} y$. Then $\rho(P^*) = \rho(P)$, and $\Delta(P^*) = * \otimes * (\tau_{\mathcal{R}} \Delta P)$, where $\tau_{\mathcal{R}}$ is a ring homomorphism $\mathcal{R} \otimes \mathcal{R} \rightarrow \mathcal{R} \otimes \mathcal{R}$ that interchanges the tensor factors $\tau_{\mathcal{R}}(x \otimes y) = y \otimes x$.

Definition 2.27. A join of polytopes defines a bilinear operation of degree $+2$ on the ring $\mathcal{P}$. It is associative and commutative, so $(\mathcal{P}, *)$ is a commutative associative ring without a unit.

Let us add a formal unit $\emptyset$ of degree -2 corresponding to the empty set. Thus $\emptyset * P = P * \emptyset = P$.

Let us denote the ring $(\mathcal{P} \oplus \mathbb{Z}\emptyset, *)$ by $\mathcal{RP}$. Then $\mathcal{RP}$ is a commutative associative ring.

Given two polytopes: P^{n_1} with v_1 vertices and m_1 facets, and Q^{n_2} with v_2 vertices and m_2 facets, the polytope $P^{n_1} * Q^{n_2}$ has dimension $n_1 + n_2 + 1$, while its number of vertices is $v_1 + v_2$, and its number of facets is $m_1 + m_2$.

Therefore $\mathcal{RP}$ is a threegraded ring with graduations $2(n+1)$, $2(v-n-1)$, and $2(m-n-1)$, where $n = \dim P$, v is the number of vertices, and m is the number of facets. The duality operator $*$ interchanges the graduations $2(v-n-1)$ and $2(m-n-1)$.

Remark 2.28. Let us recall that in the ring $\mathcal{P}$ the polytope P has two graduations $2n$ and $2(m-n)$. We have $v(P \times Q) = v(P) \cdot v(Q)$, so the number of vertices does not give the third graduation in this case.

Definition 2.29. Let us denote by $\mathcal{SP}$ the graded abelian subgroup consisting of all self-dual elements of the ring $\mathcal{RP}$. According to Proposition 2.18 it is a subring in $\mathcal{RP}$.

Example 2.30. $\mathcal{SP}$ contains the subring generated by simplices and all polygons M_n^2 .

The correspondence $P \rightarrow L(P)$ defines the homomorphism of abelian groups $L : \mathcal{P} \oplus \mathbb{Z}\emptyset \rightarrow \mathcal{R}$. It is an injection. Since $L(P * Q) = L(P) \cdot L(Q)$, it is a homomorphism of graded rings $\mathcal{RP} \rightarrow \mathcal{R}$. Moreover, $L(P^*) = L(P)^*$.

Definition 2.31. Since

$$\Delta(L(P)) = \sum_{\emptyset \subseteq F \subseteq P} [\emptyset, F] \otimes [F, P] = \sum_{\emptyset \subseteq F \subseteq P} L(F) \otimes L(P/F),$$

we see that $L(\mathcal{RP})$ is a Hopf subalgebra in $\mathcal{R}$, therefore it induces a Hopf algebra structure on the ring $\mathcal{RP}$:

- The comultiplication in $\mathcal{RP}$ is

$$\Delta(P) = \sum_{\emptyset \subseteq F \subseteq P} F \otimes P/F;$$

- The counit is

$$\varepsilon(P) = \begin{cases} 1, & \text{if } P = \emptyset, \\ 0, & \text{else;} \end{cases}$$

- The antipode χ is defined as

$$\chi(P) = \sum_{k \geq 0} \sum_{\emptyset = F_0 \subset F_1 \subset \dots \subset F_k = P} (-1)^k (F_1/F_0) * (F_2/F_1) * \dots * (F_k/F_{k-1}).$$

Remark 2.32. Let $\tau_{\mathcal{RP}} : \mathcal{RP} \otimes \mathcal{RP} \rightarrow \mathcal{RP} \otimes \mathcal{RP}$ be the ring homomorphism that interchanges the tensor factors: $\tau_{\mathcal{RP}}(x \otimes y) = y \otimes x$. Then $\Delta(P^*) = * \otimes * (\tau_{\mathcal{RP}} \Delta P)$.

Proposition 2.33. $\mathcal{RP}$ is a ring of polynomials generated by all join-indecomposable polytopes.

See Appendix B for the proof.

Definition 2.34. For the ring $\mathcal{RP}$ let us define an α -character ε_α by the formula

$$\varepsilon_\alpha(P^n) = \alpha^{n+1} = \alpha^{\rho(P)}$$

Then ε_0 is the counit ε .

Definition 2.35. Let $\mathcal{B}$ be the abelian subgroup in $\mathcal{RP}$ generated by $\emptyset$, and all the simplices $\Delta^n, n \geq 0$. Then $L(\Delta^{n-1}) = B_n$ is a Boolean algebra $\{\hat{0}, \hat{1}\}^n$. We have $\Delta^k * \Delta^l = \Delta^{k+l+1}$, so it is a subring in $\mathcal{RP}$.

Let us denote $x = \Delta^0 = \text{pt}$. Then $\Delta^{n-1} = x^n$. We have $\Delta x = 1 \otimes x + x \otimes 1$, $\chi(x) = -x$, so $\mathcal{B}$ is a Hopf subalgebra isomorphic to the Hopf algebra of polynomials $\mathbb{Z}[x]$ with the comultiplication $\Delta x = 1 \otimes x + x \otimes 1$.

Definition 2.36. We will denote by $\mathbf{R}$ the ring $\mathcal{P}$ or the ring $\mathcal{RP}$, and by $\mathbf{R}^{[2n]}$ the sum

$$\mathbf{R}^0 \oplus \mathbf{R}^2 \oplus \mathbf{R}^4 \oplus \dots \oplus \mathbf{R}^{2n}$$

2.4 Operators

Definition 2.37. Let us define the ring of linear operators $\mathcal{L}(\mathbf{R}) = \text{Hom}_{\mathbb{Z}}(\mathbf{R}, \mathbf{R})$ with the multiplication given by a composition of operators.

Definition 2.38. For $k \geq 0$ let us define *faces operators* d_k that sent a polytope P^n to the sum of all its $(n - k)$ -dimensional faces.

$$d_k P^n = \sum_{F^{n-k} \subseteq P^n} F^{n-k}$$

In particular, $d_{n+1} P^n = \begin{cases} \emptyset & \text{in } \mathcal{RP}, \\ 0 & \text{in } \mathcal{P}, \end{cases} \quad d_k \emptyset = 0, \quad k \geq 1.$

Example 2.39. We have $d_1 \text{pt} = \emptyset$ in $\mathcal{RP}$, and 0 in $\mathcal{P}$.

Definition 2.40. Let us define by $\mathcal{D}(\mathbf{R}) \subset \mathcal{L}(\mathbf{R})$ the ring of operators generated by faces operators d_k , $k \geq 0$, and by $\mathcal{D}(\mathbf{R}^{2n}, \mathbf{R}^{2(n-k)})$ the set of all operators in $\mathcal{D}(\mathbf{R})$ of degree $-2k$ acting on the abelian group $\mathbf{R}^{2n}$.

Proposition 2.41. For any two polytopes P and Q , and $d = d_1$ we have:

$$d(P \times Q) = (dP) \times Q + P \times (dQ) \text{ in } \mathcal{P}, \text{ and } d(P * Q) = (dP) * Q + P * (dQ) \text{ in } \mathcal{RP}.$$

Therefore $\mathcal{P}$ and $\mathcal{RP}$ are differential rings.

Proof. Each facet of $P \times Q$ has the form $F \times Q$ or $P \times G$, where F, G are facets of P and Q . Therefore,

$$d(P \times Q) = \sum_{F \subset P} F \times Q + \sum_{G \subset Q} P \times G = (dP) \times Q + P \times (dQ)$$

Each facet of $P * Q$ has the form $F * Q$ or $P * G$, where F, G are facets of P and Q . Therefore,

$$d(P * Q) = (dP) * Q + P * (dQ)$$

□

Example 2.42. $xP = \text{pt} * P = CP$ is a *cone*. For a polytope P of positive dimension we have

$$d(CP) = d(\text{pt} * P) = (d\text{pt}) * P + \text{pt} * (dP) = \emptyset * P + C(dP) = P + C(dP)$$

Thus, the multiplication by $x = \text{pt}$ in the ring $\mathcal{RP}$ defines a *cone operator* $C \in \mathcal{L}(\mathcal{RP})$ of degree +2 such that

$$[d, C] = \text{id}$$

On the other hand, in the ring $\mathcal{P}$ the same relation holds for all the polytopes except for $P = \text{pt}$. In this case we have $d(C\text{pt}) = dI = 2\text{pt}$, while $C(d\text{pt}) + \text{pt} = \text{pt}$. Therefore, in the ring $\mathcal{P}$ we have

$$[d, C] = \text{id} + \xi_0$$

Example 2.43. We have

$$d(\Delta^k * \Delta^l) = (k+1)\Delta^{k-1} * \Delta^l + (l+1)\Delta^k * \Delta^{l-1} = (k+l+2)\Delta^{k+l} = d\Delta^{k+l+1}$$

Example 2.44. Since $\Delta^k = x^{k+1}$, we see that $dx^{k+1} = (k+1)x^k$. Therefore $\mathbb{Z}[x]$ is a differential subring in $(\mathcal{RP}, d)$ with $d = \frac{d}{dx}$.

Definition 2.45. The correspondence $P \rightarrow BP$ defines a linear *bipyramid operator* of degree +2 on the rings $\mathcal{P}$ and $\mathcal{RP}$. By definition $B\emptyset = \text{pt}$.

Remark 2.46. It is easy to see that if P is simple and *self-dual*, i.e. P^* is combinatorially equivalent to P , then P is a simplex. There are many examples of self-dual non-simple polytopes, with simplest infinite family given by k -gonal pyramids for $k \geq 4$. The next example gives a more interesting *regular* self-dual polytope.

Example 2.47 (24-cell). Let Q be the 4-polytope obtained by taking the convex hull of the following 24 points in $\mathbb{R}^4$: endpoints of 8 vectors $\pm \mathbf{e}_i$, $1 \leq i \leq 4$, and 16 points of the form $(\pm \frac{1}{2}, \pm \frac{1}{2}, \pm \frac{1}{2}, \pm \frac{1}{2})$. It can be shown that all the facets of Q are octahedra.

The polar polytope Q^* is specified by the following 24 inequalities:

$$\pm x_i + 1 \geq 0 \text{ for } 1 \leq i \leq 4, \quad \text{and} \quad \frac{1}{2}(\pm x_1 \pm x_2 \pm x_3 \pm x_4) + 1 \geq 0. \quad (2)$$

Each of these inequalities turns into equality in exactly one of the specified 24 points, so it defines a supporting hyperplane whose intersection with Q is only one point. This implies that Q has exactly 24 vertices. The vertices of Q^* may be determined using the “elimination process” to (2) (see [Z2, Section 1.2]), and as the result we obtain 24 points of the form $\pm \mathbf{e}_i \pm \mathbf{e}_j$ for $1 \leq i < j \leq 4$. Each supporting hyperplane defined by (2) contains exactly 6 vertices of Q^* , which form an octahedron. So both Q and Q^* have 24-vertices and 24 octahedral facets. In fact, both Q and Q^* provide examples of a *regular 4-polytope* called a *24-cell*. It is the only regular self-dual polytope different from a simplex. For more details on 24-cell and other regular polytopes see [C].

In this example we have $dQ = 24BI^2$, and $(dQ)^* = 24I^3$. Therefore for a self-dual element $P \in \mathcal{SP}$, dP can be non self-dual. So $\mathcal{SP}$ is not a differential subring in $\mathcal{RP}$.

However there is another derivation defined on $\mathcal{SP}$. Since the duality $*$ is a ring homomorphism, $\delta = *d*$ is a derivation of the ring $\mathcal{RP}$, as well as d .

Proposition 2.48. $d + \delta$ is a derivation of the ring $\mathcal{SP}$

Proof. Since for any $P \in \mathcal{SP}$, $P^* = P$, we have

$$((d + \delta)P)^* = *(d + *d*)P = *dP + **dP = *d * P + dP = (\delta + d)P.$$

□

Proposition 2.49.

$$\begin{aligned} d_k(P \times Q) &= \sum_{i+j=k} (d_i P) \times (d_j Q) \quad \text{in } \mathcal{P} \\ d_k(P * Q) &= \sum_{i+j=k} (d_i P) * (d_j Q) \quad \text{in } \mathcal{RP}. \end{aligned}$$

Proof. We have

$$\begin{aligned} d_k(P^n \times Q^r) &= \sum_{F^{n-i} \subseteq P^n, G^{r-j} \subseteq Q^r, i+j=k} F^{n-i} \times G^{r-j} = \sum_{i+j=k} \sum_{F^{n-i} \subseteq P^n} F^{n-i} \times \left(\sum_{G^{r-j} \subseteq Q^r} G^{r-j} \right) = \\ &= \sum_{i+j=k} \left(\sum_{F^{n-i} \subseteq P^n} F^{n-i} \right) \times \left(\sum_{G^{r-j} \subseteq Q^r} G^{r-j} \right) = \sum_{i+j=k} (d_i P) \times (d_j Q) \end{aligned}$$

The same argument works also for $P^n * Q^r$. The only difference is that faces of the form $\emptyset * G^{r-j}$, $i = n+1, i+j = k$, and $F^{n-i} * \emptyset$, $j = r+1, i+j = k$ are allowed. Here $\emptyset = d^{n+1}P^n = d^{r+1}Q^r$. □

Definition 2.50. Let us define an operator $\Phi : \mathcal{R} \rightarrow \mathcal{R}[t]$, $\Phi(t) \in \mathcal{D}(\mathcal{R})[[t]]$ by the formula

$$\Phi(t) = 1 + dt + d_2 t^2 + \dots + d_k t^k + \dots$$

Definition 2.51. Let $f_i(P^n)$ be the number of i -dimensional faces of P^n .

Proposition 2.52. We have $\Phi(-t)\Phi(t) = 1$ on the ring $\mathcal{R}$, that is for any $n \geq 1$

$$d_n - dd_{n-1} + \dots + (-1)^{n-1} d_{n-1} d + (-1)^n d_n = 0.$$

Proof. The relation is clear for $\emptyset \in \mathcal{RP}$ and for $\text{pt} \in \mathcal{P}$. For $\text{pt} \in \mathcal{RP}$ we have

$$\Phi(-t)\Phi(t) \text{pt} = (1 - dt + \dots)(\text{pt} + \emptyset t) = \text{pt} - \emptyset t + \emptyset t = \text{pt}$$

Let us consider the polytope P^n of positive dimension.

$$\Phi(-t)\Phi(t) = \sum_{k=0}^{\infty} \left(\sum_{i=0}^k (-1)^i d_i d_{k-i} \right) t^k.$$

If $k > n+1$ then the coefficient of t^k in the series $\Phi(-t)\Phi(t)P^n$ is equal to 0. If $k = 0$, then it is equal to P^n . Let $1 \leq k \leq n$. Then

$$\begin{aligned} (dd_{k-1} - d_2 d_{k-2} + \dots + (-1)^{k-2} d_{k-1} d) P^n &= \\ &= \sum_{F^{n-k} \subset P^n} \left(\sum_{F^{n-k+1} \supset F^{n-k}} 1 - \sum_{F^{n-k+2} \supset F^{n-k}} 1 + \dots + (-1)^{k-2} \sum_{F^{n-1} \supset F^{n-k}} 1 \right) F^{n-k} = \\ &= \sum_{F^{n-k} \subset P^n} (f_0(P^n/F^{n-k}) - f_1(P^n/F^{n-k}) + \dots + (-1)^{k-2} f_{k-2}(P^n/F^{n-k})) F^{n-k} \end{aligned}$$

Since P^n/F^{n-k} is a $(k-1)$ -dimensional polytope, the Euler formula gives the relation

$$f_0(P^n/F^{n-k}) - f_1(P^n/F^{n-k}) + \dots + (-1)^{k-2} f_{k-2}(P^n/F^{n-k}) = 1 + (-1)^k.$$

Thus we obtain

$$(dd_{k-1} - d_2d_{k-2} + \cdots + (-1)^{k-2}d_{k-1}d)P^n = (1 + (-1)^k)d_kP^n \Leftrightarrow \sum_{i=0}^k (-1)^i d_i d_{k-i} P^n = 0$$

The coefficient of t^{n+1} is equal to 0 in $\mathcal{P}$, and to $(1 - f_0 + f_1 + \cdots + (-1)^n f_{n-1} + (-1)^{n+1})\emptyset$ in $\mathcal{RP}$. Then the Euler formula implies that it is equal to 0. $\square$

Proposition 2.53.

$$\begin{aligned} \xi_{-\alpha}\Phi(\alpha) &= \xi_\alpha \quad \text{in the ring } \mathcal{P} \\ \varepsilon_{-\alpha}\Phi(\alpha) &= \varepsilon_0 \quad \text{in the ring } \mathcal{RP}. \end{aligned}$$

Proof. Let us note that $\xi_\alpha d_k P^n = f_{n-k}(P^n)\alpha^{n-k}$. Then for any polytope P^n we have

$$\xi_{-\alpha}\Phi(\alpha)P^n = (-\alpha)^n + (-\alpha)^{n-1}f_{n-1}\alpha + \cdots + f_0\alpha^n = \left(\sum_{i=0}^n (-1)^i f_i\right)\alpha^n = \alpha^n = \xi_\alpha P^n,$$

since the Euler formula gives the relation:

$$f_0 - f_1 + \cdots + (-1)^{n-1}f_{n-1} + (-1)^n = 1$$

On the other hand, in the ring $\mathcal{RP}$ we have $\varepsilon_\alpha d_k P^n = f_{n-k}(P^n)\alpha^{n-k+1}$, so

$$\varepsilon_{-\alpha}\Phi(\alpha)P^n = (-\alpha)^{n+1} + (-\alpha)^n f_{n-1}\alpha + \cdots + (-\alpha)f_0\alpha^n + \alpha^{n+1}$$

According to the Euler formula this expression is equal to 0 for all the polytopes but $\emptyset$. Thus $\varepsilon_{-\alpha}\Phi(\alpha) = \varepsilon_0$. $\square$

Example 2.54. On the ring of simple polytopes $\mathcal{P}_s \subset \mathcal{P}$ we have the relations $k!d_k|_{\mathcal{P}_s} = d^k|_{\mathcal{P}_s}$, therefore $\mathcal{D}(\mathcal{P}_s)$ is isomorphic to the divided power ring $d_k d_l = \binom{k+l}{k} d_{k+l}$, and $\Phi(t)|_{\mathcal{P}_s} = e^{dt}$.

Let us note that this equality is not valid for $\mathcal{P}_s \subset \mathcal{RP}$. The problem is that for a simple polytope P^n we have $d_k = \frac{d^k}{k!}$ for all $1 \leq k \leq n$. But $d_{n+1}P^n = \emptyset$, while

$$\frac{d^{n+1}}{(n+1)!}P^n = \frac{d}{n+1} \left(\frac{d^n}{n!}P^n \right) = \frac{d}{n+1} (d_n P^n) = \frac{d}{n+1} f_0(P^n) \text{pt} = \frac{f_0(P^n)}{n+1} \emptyset$$

Another example of operators in $\mathcal{L}(\mathbf{R})$ is given by the multiplication by elements of $\mathbf{R}$:

$$[P](Q) = PQ$$

Proposition 2.55. *The following relation holds*

$$d_k[P] = \sum_{i=0}^k [d_i P] d_{k-i}$$

Proof.

$$d_k(PQ) = \sum_{i=0}^k (d_i P) d_{k-i} Q$$

$\square$

Thus any operator in the ring $\mathbf{R}\mathcal{D}(\mathbf{R})$ generated by $\mathcal{D}(\mathbf{R})$ and $\{[P], P \in \mathbf{R}\}$ can be expressed as a sum of operators

$$\sum_{P, \omega} [P] D_\omega, \quad P \in \mathbf{R}, \quad D_\omega \in \mathcal{D}(\mathbf{R}).$$

Let us remind that $[\text{pt}] = x$ is a cone operator in $\mathcal{L}(\mathcal{RP})$.

2.5 Flag f -vectors

Definition 2.56. Let P^n be an n -dimensional polytope and $S = \{a_1, \dots, a_k\} \subset \{0, 1, \dots, n-1\}$. A *flag number* $f_S = f_{a_1, \dots, a_k}$ is the number of increasing sequences of faces

$$F^{a_1} \subset F^{a_2} \subset \dots \subset F^{a_k}, \quad \dim F^{a_i} = a_i.$$

It is easy to see that for $S = \{i\}$ the number $f_{\{i\}} = f_i$ is just the number of i -dimensional faces. We have already defined this number above. Let us denote $l(S) = k$. Then $0 \leq k \leq n$.

The collection $\{f_S\}$ of all the flag numbers is called a *flag f -vector* (or *extended f -vector*) of the polytope P^n . By the definition $f_\emptyset = 1$.

Remark 2.57. For the set $\{a_1, \dots, a_k\} \subset \{0, 1, \dots, n-1\}$ let us denote $a_0 = -1$, and $a_{k+1} = n$. Then

$$f_{-1, a_1, \dots, a_k} = f_{-1, a_1, \dots, a_k, n} = f_{a_1, \dots, a_k, n} = f_{a_1, \dots, a_k}$$

This notation corresponds to the fact that $\emptyset$ is the only face of P of dimension -1 , and any other face contains $\emptyset$, while the polytope P itself is the only face of dimension n , and it contains any other face.

Flag f -vectors have been extensively studied by M. Bayer and L. Billera in [BB], where the generalized Dehn-Sommerville relations were proved:

Theorem ([BB], Theorem 2.1). *Let P^n be an n -dimensional polytope, and $S \subset \{0, \dots, n-1\}$. If $\{i, k\} \subseteq S \cup \{-1, n\}$ such that $i < k-1$ and $S \cap \{i+1, \dots, k-1\} = \emptyset$, then*

$$\sum_{j=i+1}^{k-1} (-1)^{j-i-1} f_{S \cup \{j\}} = (1 - (-1)^{k-i-1}) f_S. \quad (3)$$

Definition 2.58. For $n \geq 1$ let Ψ^n be the set of subsets $S \subset \{0, 1, \dots, n-2\}$ such that S contains no two consecutive integers.

It is easy to show by induction that the cardinality of Ψ^n is equal to the n -th Fibonacci number c_n ($c_n = c_{n-1} + c_{n-2}$, $c_0 = 1$, $c_1 = 1$).

Let us remind that for any polytope P we have defined a *cone* (or a *pyramid*) CP and a *bipyramid* (or a *suspension*) BP . These two operations are defined on combinatorial polytopes and can be extended to linear operators on the rings $\mathcal{P}$ and $\mathcal{RP}$. By definition $B\emptyset = \text{pt} = C\emptyset$.

The face lattice of the polytope CP^n is:

$$\{\emptyset\}; \{F_i^0, C\emptyset\}; \{F_i^1, CF_j^0\}; \dots; \{F_i^{n-1}, CF_j^{n-2}\}; \{P^n, CF_i^{n-1}\}; \{CP^n\},$$

where F_i^k are k -dimensional faces of P^n , and

$$\begin{aligned} F^{a_1} \subset F^{a_2} \text{ in } CP &\Leftrightarrow F^{a_1} \subset F^{a_2} \text{ in } P, \\ F^{a_1} \subset CF^{a_2} &\Leftrightarrow F^{a_1} \subset F^{a_2}, \\ CF^{a_1} \subset CF^{a_2} &\Leftrightarrow F^{a_1} \subset F^{a_2}. \end{aligned}$$

The polytope BP^n has the face lattice:

$$\{\emptyset\}, \{C_{-1}\emptyset, C_1\emptyset, F_i^0\}, \{F_i^1, C_{-1}F_j^0, C_1F_k^0\}, \dots, \{F_k^{n-1}, C_{-1}F_j^{n-2}, C_1F_k^{n-2}\}, \{C_{-1}F_i^{n-1}, C_1F_j^{n-1}\}, \{BP^n\}.$$

where C_{-1} and C_1 are the lower and the upper cones, and

$$\begin{aligned} F^{a_1} \subset F^{a_2} \text{ in } BP &\Leftrightarrow F^{a_1} \subset F^{a_2} \text{ in } P, \\ F^{a_1} \subset C_s F^{a_2} &\Leftrightarrow F^{a_1} \subset F^{a_2}; \\ C_s F^{a_1} \subset C_t F^{a_2} &\Leftrightarrow s = t \text{ and } F^{a_1} \subset F^{a_2}; \end{aligned}$$

Definition 2.59. For $n \geq 1$ let Ω^n be the set of n -dimensional polytopes that arises when we apply words in B and C that end in C^2 and contain no adjacent B 's to the empty set $\emptyset$.

Each word of length $n+1$ from the set Ω^n either has the form CQ , $Q \in \Omega^{n-1}$, or BCQ , $Q \in \Omega^{n-2}$, so cardinality of the set Ω^n satisfies the Fibonacci relation $|\Omega^n| = |\Omega^{n-1}| + |\Omega^{n-2}|$. Since $|\Omega^1| = |\{C^2\}| = 1$, and $|\Omega^2| = |\{C^3, BC^2\}| = 2$, we see that $|\Omega^n| = c_n = |\Psi^n|$.

M. Bayer and L. Billera proved the following fact:

Theorem. *Let $n \geq 1$. Then*

1. *For all $T \subseteq \{0, 1, \dots, n-1\}$ there is a nontrivial linear relation expressing $f_T(P)$ in terms of $f_S(P)$, $S \in \Psi^n$, which holds for all n -dimensional polytopes (see [BB, Proposition 2.2]).*
2. *The extended f -vectors of the c_n elements of Ω^n are affinely independent [BB, Proposition 2.3]. Thus the flag f -vectors*

$$\{f_S(P^n)\}_{S \in \Psi^n} : P^n - \text{an } n\text{-dimensional polytope}$$

span an $(c_n - 1)$ -dimensional affine hyperplane defined by the equation $f_\emptyset = 1$ (see [BB, Theorem 2.6])

Remark 2.60. The first part of the theorem follows from the generalized Dehn-Sommerville relations (3). In fact, we have $f_{S, n-2, n-1} = 2f_{S, n-2}$, and

$$f_{S, i, n-1} = (1 + (-1)^{n-i})f_{S, i} + (-1)^{n-i-1} \sum_{j=i+1}^{n-2} (-1)^{j-i-1} f_{S, i, j}, \quad i < n-2.$$

Therefore, we can get rid of the index $n-1$. Then $f_{S_1, i-1, i, i+1, S_2} = 2f_{S_1, i-1, i+1, S_2}$. Therefore, we can get rid of triples of consecutive indices.

Now for each set S in the sum we obtain let us take the last pair of consecutive indices $\{i, i+1\}$. Then $f_S = f_{S_1, j, i, i+1, S_2}$.

$$f_{S_1, j, i, i+1, S_2} = (1 + (-1)^{i+1-j})f_{S_1, j, i+1, S_2} + (-1)^{i-j} \sum_{k=j+1}^{i-1} (-1)^{k-j-1} f_{S_1, j, k, i+1, S_2}$$

Then the last pair of consecutive indices for each set in the sum consists of smaller numbers. Then after the sequence of such steps and removing the triples of consecutive indices on each step we will obtain flag numbers of the form $f_{i, i+1, S_2}$, where the set $\{i+1\} \cup S_2$ contains no consecutive indices. Then

$$f_{i, i+1, S_2} = (1 + (-1)^i)f_{i+1, S_2} - (-1)^i \sum_{j=0}^{i-1} (-1)^j f_{j, i+1, S_2}$$

where all the sets $\{j, i+1\} \cup S_2$ belong to Ψ^n . This finishes the proof.

In fact, a little bit more stronger statement, which can be easily extracted from the original Bayer-Billera's proof, is true.

Let us identify the words in Ω^n with the sets in Ψ^n in such a way that the word $C^{n+1-a_k}BC^{a_k-a_{k-1}-1}B \dots BC^{a_1-1}$ corresponds to the set $\{a_1-3, \dots, a_k-3\}$. Let us set $C < B$ and order the words lexicographically. Consider a matrix K^n of sizes $c_n \times c_n$, $k_{Q, S} = f_S(Q)$, $Q \in \Omega^n$, $S \in \Psi^n$.

Proposition 2.61.

$$\det(K^n) = 1.$$

Proof. For $n = 1$ the matrix $K^1 = (f_\emptyset(I)) = (1)$.

For $n = 2$

$$K^2 = \begin{pmatrix} 1 & f_0(\Delta^2) \\ 1 & f_0(I^2) \end{pmatrix} = \begin{pmatrix} 1 & 3 \\ 1 & 4 \end{pmatrix}; \quad \det K^2 = 1$$

Let us prove the statement by induction. Consider the matrix K^n :

$$K^n = \begin{pmatrix} K_{11} & K_{12} \\ K_{21} & K_{22} \end{pmatrix}$$

where the block K_{11} corresponds to the words of the form CQ , $Q \in \Omega^{n-1}$ and the sets S , that do not contain $n-2$. Similarly, the block K_{22} corresponds to the words of the form BCQ , $Q \in \Omega^{n-2}$ and the sets containing $n-2$.

Each increasing sequence of faces of the polytope CP^{n-1} without two faces of adjacent dimensions has the form

$$F^{l_1} \subset \dots \subset F^{l_i} \subset CF^{l_{i+1}} \subset \dots \subset CF^{l_k},$$

so for $\{a_1, \dots, a_k\} \in \Psi^n$

$$f_{a_1, \dots, a_k}(CP^{n-1}) = f_{a_1-1, a_2-1, \dots, a_k-1}(P^n) + \dots + f_{a_1, \dots, a_i, a_{i+1}-1, \dots, a_k-1}(P^n) + \dots + f_{a_1, \dots, a_k}(P^n), \quad (4)$$

where $f_{-1, a_2, \dots, a_k} = f_{a_2, \dots, a_k}$.

The sets of the form $\{a_1, \dots, a_l, a_{l+1}-1, \dots, a_k-1\}$ may not belong to Ψ^{n-1} . But we can express the corresponding flag numbers in terms of $\{f_S, S \in \Psi^{n-1}\}$ using the generalized Dehn-Sommerville relations. Let $a_k < n-2$. Each relation has the form:

$$f_{S_1, i, k-1, k, S_2} = (-1)^{k-i} \left((1 - (-1)^{k-i-1}) f_{S_1, i, k, S_2} - \sum_{j=i+1}^{k-2} (-1)^{j-i-1} f_{S_1, i, j, k, S_2} \right),$$

where all the sets on the right side are lower than the set on the left side. Thus for $a_k < n-2$

$$f_{a_1, \dots, a_k}(CP^n) = f_{a_1, \dots, a_k}(P^n) + \text{lower summands},$$

so the matrix K_{11} can be represented as $K^{n-1}T$, where T is an upper unitriangular matrix. In particular, $\det K_{11} = \det K^{n-1} \cdot \det T = 1$ by induction.

Lemma 2.62. *Let P be an $(n-2)$ -dimensional polytope, and $S \in \Psi^n$. Then for the polytopes BCP and CBP*

$$f_S(BCP) = \begin{cases} f_S(CBP), & \text{if } n-2 \notin S, \\ f_S(CBP) + f_{S \setminus \{n-2\}}(P), & \text{if } n-2 \in S, \end{cases}$$

Proof. The polytope BCP has faces

$$\{F, \emptyset \subseteq F \subseteq P\}; \{CF, \emptyset \subseteq F \subsetneq P\}; \{C_a F, \emptyset \subseteq F \subseteq P\}; \{C_a CF, \emptyset \subseteq F \subsetneq P\}; BCP,$$

where $a = \pm 1$

Then each increasing sequence of faces corresponding to the set S has one of the forms

- $F^{l_1} \subset \dots \subset F^{l_i}$
- $F^{l_1} \subset \dots \subset F^{l_i} \subset CF^{l_{i+1}} \subset \dots \subset CF^{l_j}$,
- $F^{l_1} \subset \dots \subset F^{l_i} \subset C_a F^{l_{i+1}} \subset \dots \subset C_a F^{l_j}$
- $F^{l_1} \subset \dots \subset F^{l_i} \subset CF^{l_{i+1}} \subset \dots \subset CF^{l_j} \subset C_a CF^{l_{j+1}} \subset \dots \subset C_a CF^{l_k}$
- $F^{l_1} \subset \dots \subset F^{l_i} \subset C_a F^{l_{i+1}} \subset \dots \subset C_a F^{l_j} \subset C_a CF^{l_{j+1}} \subset \dots \subset C_a CF^{l_k}$
- $F^{l_1} \subset \dots \subset F^{l_j} \subset C_a CF^{l_{j+1}} \subset \dots \subset C_a CF^{l_k}$

The polytope CBP has faces

$$\{F, \emptyset \subseteq F \subsetneq P\}; \{C_a F, \emptyset \subseteq F \subsetneq P\}; BP; \{CF, \emptyset \subseteq F \subsetneq P\}; \{C C_a F, \emptyset \subseteq F \subsetneq P\}; CBP,$$

where $a = \pm 1$

Then each increasing sequence of faces corresponding to the set S has one of the forms

- $F^{l_1} \subset \dots \subset F^{l_i}, F^{l_i} \neq P$,
- $F^{l_1} \subset \dots \subset F^{l_i} \subset CF^{l_{i+1}} \subset \dots \subset CF^{l_j}$,
- $F^{l_1} \subset \dots \subset F^{l_i} \subset C_a F^{l_{i+1}} \subset \dots \subset C_a F^{l_j}$

- $F^{l_1} \subset \dots \subset F^{l_i} \subset CF^{l_{i+1}} \subset \dots \subset CF^{l_j} \subset CC_a F^{l_{j+1}} \subset \dots \subset C_a C F^{l_k}$
- $F^{l_1} \subset \dots \subset F^{l_i} \subset C_a F^{l_{i+1}} \subset \dots \subset C_a F^{l_j} \subset CC_a F^{l_{j+1}} \subset \dots \subset CC_a F^{l_k}$
- $F^{l_1} \subset \dots \subset F^{l_j} \subset CC_a F^{l_{j+1}} \subset \dots \subset CC_a F^{l_k}$

We can interchange CC_a and $C_a C$ to obtain a one-to-one correspondence between the increasing sequences containing both C and C_a . There is a natural bijection between sequences containing exactly one of these operations, as well as between sequences containing neither operations nor P . So $f_S(BCP) = f_S(CBP)$, if $n-2 \notin S$.

The only difference appears in the case when the sequence of faces in the polytope BCP has the form

$$F^{a_1} \subset \dots \subset F^{a_{k-1}} \subset P$$

The sequences of this type give exactly $f_{a_1, \dots, a_{k-1}}(P)$, so $f_S(BCP) = f_S(CBP) + f_{S \setminus \{n-2\}}(P)$, if $n-2 \in S$. $\square$

Let us consider the polytope BCQ corresponding to one of the lower rows of K^n . If Q starts with C , then there is a row CBQ in the upper part of the matrix. Let us subtract the row CBQ from the row BCQ . Then Lemma 2.62 implies that the resulting row is

$$k'_{BCQ, S} = \begin{cases} 0, & \text{if } n-2 \notin S; \\ f_{S \setminus \{n-2\}}(Q), & \text{if } n-2 \in S. \end{cases} \quad (5)$$

If Q starts with B , then there is no row CBQ in the matrix. But since $\det K^{n-1} = 1$ and all flag f -numbers of $(n-1)$ -dimensional polytopes can be expressed in terms of $\{f_S, S \in \Psi^{n-1}\}$, the flag f -vector of any $(n-1)$ -dimensional polytope is an integer combination of flag f -vectors of the polytopes from Ω^{n-1} . So the flag f -vector of the polytope BQ can be expressed as an integer combination of flag f -vectors of the polytopes $Q' \in \Omega^{n-1}$:

$$f(BQ) = \sum_{Q'} n_{Q'} f(Q')$$

Using Formula (4) we obtain:

$$f(CBQ) = \sum_{Q'} n_{Q'} f(CQ')$$

If we subtract the corresponding integer combination of the rows of the upper part of the matrix from the row BCQ , we obtain the row (5).

Thus we see, that using elementary transformations of rows the matrix K^{n-1} can be transformed to the matrix

$$\begin{pmatrix} K^{n-1T} & K_{12} \\ 0 & K^{n-2} \end{pmatrix}$$

By the inductive assumption $\det K^{n-1} = \det K^{n-2} = 1$, so $\det K^n = 1$. $\square$

Remark 2.63. In the proof we follow the original Bayer and Billera's idea, except for the fact that they use the additional matrix of face numbers of the polytopes $Q \in \Psi^n$, and our proof is direct.

Corollary 2.64. *The flag f -vector of any n -dimensional polytope P^n is an integer combination of the flag f -vectors of the polytopes $Q \in \Omega^n$.*

Proof. Indeed, any flag f -number is a linear combination of $f_S, S \in \Psi^n$. Since $\det K^n = 1$, the vector $\{f_S(P), S \in \Psi^n\}$ is an integer combination of the vectors $\{f_S(Q), S \in \Psi^n\}, Q \in \Omega^n$.

This implies that the whole vector $\{f_S(P)\}$ is an integer combination of the vectors $\{f_S(Q)\}, Q \in \Omega^n$ with the same coefficients. $\square$

This fact is important, when we try to describe the image of the generalized f -polynomial in the ring of quasi-symmetric functions (see below).

Generalized f -vectors are also connected with a very interesting construction of a cd -index invented by J. Fine (see [BK, Prop. 2], see also the papers by R. Stanley [St3] and M. Bayer, A. Klapper [BK]).

3 Hopf algebras

In this part we follow mainly the notations of [H]. See also [BR] and [CFL].

3.1 Quasi-symmetric functions

Definition 3.1. A *composition* ω of a number n is an ordered set $\omega = (j_1, \dots, j_k)$, $j_i \geq 1$, such that $n = j_1 + \dots + j_k$. Let us denote $|\omega| = n$, $l(\omega) = k$.

Let us denote by $()$ the empty composition of the number 0. Then $|\emptyset| = 0$, $l(\emptyset) = 0$.

Definition 3.2. Let $t_1, t_2, \dots$ be a finite or an infinite set of variables, $\deg t_i = 2$. For a composition $\omega = (j_1, \dots, j_k)$ consider a *quasi-symmetric monomial*

$$M_\omega = \sum_{l_1 < \dots < l_k} t_{l_1}^{j_1} \dots t_{l_k}^{j_k}, \quad M_{\emptyset} = 1.$$

Degree of the monomial M_ω is equal to $2|\omega| = 2(j_1 + \dots + j_k)$.

For any two monomials $M_{\omega'}$ and $M_{\omega''}$ their product in the ring of polynomials $\mathbb{Z}[t_1, t_2, \dots]$ is equal to

$$M_{\omega'} M_{\omega''} = \sum_{\omega} \left(\sum_{\Omega' + \Omega'' = \omega} 1 \right) M_\omega,$$

where for the compositions $\omega = (j_1, \dots, j_k)$, $\omega' = (j'_1, \dots, j'_{l'})$, $\omega'' = (j''_1, \dots, j''_{l''})$, Ω' and Ω'' are all the k -tuples such that

$$\Omega' = (0, \dots, j'_1, \dots, 0, \dots, j'_{l'}, \dots, 0), \quad \Omega'' = (0, \dots, j''_1, \dots, 0, \dots, j''_{l''}, \dots, 0),$$

This multiplication rule of compositions is called *the overlapping shuffle multiplication*. For example,

1.

$$M_{(1)} M_{(1)} = \left(\sum_i t_i \right) \left(\sum_j t_j \right) = \sum_i t_i^2 + 2 \sum_{i < j} t_i t_j = M_{(2)} + 2M_{(1,1)}.$$

This corresponds to the decompositions

$$(2) = (1) + (1), \quad (1, 1) = (1, 0) + (0, 1) = (0, 1) + (1, 0).$$

2.

$$\begin{aligned} M_{(1)} M_{(1,1)} &= \left(\sum_i t_i \right) \left(\sum_{j < k} t_j t_k \right) = \sum_{i < j} t_i^2 t_j + \sum_{i < j} t_i t_j^2 + 3 \sum_{i < j < k} t_i t_j t_k = \\ &= M_{(2,1)} + M_{(1,2)} + 3M_{(1,1,1)}. \end{aligned}$$

This corresponds to the decompositions

$$\begin{aligned} (2, 1) &= (1, 0) + (1, 1), \quad (1, 2) = (1, 0) + (1, 1), \\ (1, 1, 1) &= (1, 0, 0) + (0, 1, 1) = (0, 1, 0) + (1, 0, 1) = (0, 0, 1) + (1, 1, 0). \end{aligned}$$

3.

$$\begin{aligned} M_{(1,1)} M_{(1,1)} &= \left(\sum_{i < j} t_i t_j \right) \left(\sum_{k < l} t_k t_l \right) = \sum_{i < j} t_i^2 t_j^2 + 2 \sum_{i < j < k} t_i^2 t_j t_k + \\ &\quad + 2 \sum_{i < j < k} t_i t_j^2 t_k + 2 \sum_{i < j < k} t_i t_j t_k^2 + 6 \sum_{i < j < k < l} t_i t_j t_k t_l = \\ &= M_{(2,2)} + 2M_{(2,1,1)} + 2M_{(1,2,1)} + 2M_{(1,1,2)} + 6M_{(1,1,1,1)}. \end{aligned}$$

This corresponds to the decompositions

$$\begin{aligned}
(2, 2) &= (1, 1) + (1, 1), \quad (2, 1, 1) = (1, 1, 0) + (1, 0, 1) = (1, 0, 1) + (1, 1, 0), \\
(1, 2, 1) &= (1, 1, 0) + (0, 1, 1) = (0, 1, 1) + (1, 1, 0), \quad (1, 1, 2) = (1, 0, 1) + (0, 1, 1) = (0, 1, 1) + (1, 0, 1), \\
(1, 1, 1, 1) &= (1, 1, 0, 0) + (0, 0, 1, 1) = (1, 0, 1, 0) + (0, 1, 0, 1) = (1, 0, 0, 1) + (0, 1, 1, 0) = \\
&= (0, 1, 1, 0) + (1, 0, 0, 1) = (0, 1, 0, 1) + (1, 0, 1, 0) = (0, 0, 1, 1) + (1, 1, 0, 0).
\end{aligned}$$

Thus finite integer combinations of quasi-symmetric monomials form a ring. This ring is called a *ring of quasi-symmetric functions* and is denoted by $\mathcal{Qsym}[t_1, \dots, t_n]$, where n is the number of variables. In the case of an infinite number of variables it is denoted by $\mathcal{Qsym}[t_1, t_2, \dots]$ or $\mathcal{Qsym}$.

The diagonal mapping $\Delta: \mathcal{Qsym} \rightarrow \mathcal{Qsym} \otimes \mathcal{Qsym}$

$$\Delta M_{(a_1, \dots, a_k)} = \sum_{i=0}^k M_{(a_1, \dots, a_i)} \otimes M_{(a_{i+1}, \dots, a_k)}$$

defines on $\mathcal{Qsym}$ the structure of a graded Hopf algebra.

Proposition 3.3. *A polynomial $g \in \mathbb{Z}[t_1, \dots, t_r]$ is a finite linear combination of quasi-symmetric monomials if and only if*

$$g(0, t_1, t_2, \dots, t_{r-1}) = g(t_1, 0, t_2, \dots, t_{r-1}) = \dots = g(t_1, \dots, t_{r-1}, 0)$$

Proof. For the quasi-symmetric monomial M_ω we have

$$M_\omega(t_1, \dots, t_i, 0, t_{i+1}, \dots, t_{r-1}) = \begin{cases} 0, & \omega = (j_1, \dots, j_r); \\ M_\omega(t_1, \dots, t_i, t_{i+1}, \dots, t_{r-1}), & \omega = (j_1, \dots, j_k), \quad k < r. \end{cases}$$

So this property is true for all quasi-symmetric functions.

On the other hand, let the condition of the proposition be true. Let us prove that for a fixed composition ω any two monomials $t_{l_1}^{j_1} \dots t_{l_k}^{j_k}$ and $t_{l'_1}^{j_1} \dots t_{l'_k}^{j_k}$ have the equal coefficients $g_{l_1, \dots, l_k}^{j_1, \dots, j_k}$ and $g_{l'_1, \dots, l'_k}^{j_1, \dots, j_k}$.

Let $l_i + 1 < l_{i+1}$ or $i = k$ and $l_i < r$. Consider the corresponding coefficients in the polynomial equation:

$$g(t_1, \dots, t_{l_i-1}, 0, t_{l_i}, t_{l_i+1}, \dots, t_{r-1}) = g(t_1, \dots, t_{l_i-1}, t_{l_i}, 0, t_{l_i+1}, \dots, t_{r-1}).$$

On the left the monomial $t_{l_1}^{j_1} \dots t_{l_i}^{j_i} t_{l_{i+1}-1}^{j_{i+1}} \dots t_{l_k-1}^{j_k}$ has the coefficient $g_{l_1, \dots, l_i, l_{i+1}, \dots, l_k}^{j_1, \dots, j_k}$, and on the right $g_{l_1, \dots, l_i, l_{i+1}, \dots, l_k}^{j_1, \dots, j_k}$, so they are equal. Now we can move the index l_i to the right to $l_i + 1$ and in the same manner we can move l_i to the left, if $l_{i-1} < l_i - 1$, or $i = 1$ and $l_i > 1$.

Now let us move step by step the index l_1 to 1, then the index l_2 to 2, and so on. At last we obtain that $g_{l_1, \dots, l_k}^{j_1, \dots, j_k} = g_{1, \dots, k}^{j_1, \dots, j_k}$. $\square$

Remark 3.4. Similar argument shows that the same is true in the case of an infinite number of variables, if we consider all the expressions

$$a + \sum_{k=1}^{\infty} \sum_{1 \leq l_1 < \dots < l_k} \sum_{j_1, \dots, j_k \geq 1} a_{l_1, \dots, l_k}^{j_1, \dots, j_k} t_{l_1}^{j_1} \dots t_{l_k}^{j_k}.$$

of bounded degree: $2|\omega| = 2(j_1 + \dots + j_k) < N$ for all ω . Then the infinite series g of bounded degree belongs to $\mathcal{Qsym}$ if and only if

$$g(t_1, \dots, t_{i-1}, 0, t_{i+1}, \dots) = g(t_1, \dots, t_{i-1}, t_{i+1}, \dots)$$

for all $i \geq 1$. As a corollary we obtain another proof of the fact that quasi-symmetric functions form a ring.

In [H] M. Hazewinkel proved the Ditters conjecture that $\text{Qsym}[t_1, t_2, \dots]$ is a free commutative algebra of polynomials over the integers.

The $(2n)$ -th graded component $\text{Qsym}^{2n}[t_1, t_2, \dots]$ has rank 2^{n-1} .

The numbers β_i of the multiplicative generators of degree $2i$ can be found by a recursive relation:

$$\frac{1-t}{1-2t} = \prod_{i=1}^{\infty} \frac{1}{(1-t^i)^{\beta_i}}$$

There are group homomorphisms:

$$\begin{aligned} V_{r+1} : \text{Qsym}[t_1, \dots, t_r] &\rightarrow \text{Qsym}[t_1, \dots, t_{r+1}] : & V_{r+1}M_\omega(t_1, \dots, t_r) &= M_\omega(t_1, \dots, t_{r+1}) \\ E_r : \text{Qsym}[t_1, \dots, t_{r+1}] &\rightarrow \text{Qsym}[t_1, \dots, t_r] : & t_{r+1} &\rightarrow 0; \end{aligned}$$

The mapping E_r sends all the monomials M_ω corresponding to the compositions $(j_1, \dots, j_{r+1})$ of length $r+1$ to zero, and for the compositions ω of smaller length $E_r M_\omega(t_1, \dots, t_{r+1}) = M_\omega(t_1, \dots, t_r)$. It is a ring homomorphism.

It is easy to see that $E_r V_{r+1}$ is the identity map of the ring $\text{Qsym}[t_1, \dots, t_r]$.

Given $r > 0$ there is a projection $\Pi_{\text{Qsym}} : \mathbb{Z}[t_1, \dots, t_r] \rightarrow \text{Qsym}[t_1, \dots, t_r] \otimes \mathbb{Q}$:
for $s_1 < s_2 < \dots < s_k$

$$\Pi_{\text{Qsym}} t_{s_1}^{j_1} \dots t_{s_k}^{j_k} = \frac{1}{\binom{r}{k}} \left(\sum_{l_1 < \dots < l_k} t_{l_1}^{j_1} \dots t_{l_k}^{j_k} \right),$$

which gives the average value over all the monomials of each type.

Then for the quasi-symmetric monomial M_ω , $\omega = (j_1, \dots, j_k)$ we have:

$$\Pi_{\text{Qsym}} M_\omega = \sum_{1 \leq j_1 < \dots < j_k \leq r} \frac{1}{\binom{r}{k}} M_\omega = \binom{r}{k} \frac{1}{\binom{r}{k}} M_\omega = M_\omega,$$

So Π_{Qsym} is indeed a projection.

Remark 3.5. We see that in the theory of symmetric and quasi-symmetric functions there is an important additional graduation – the number of variables in the polynomial.

Definition 3.6. For a composition $\omega = (j_1, \dots, j_k)$ let us define the composition $\omega^* = (j_k, \dots, j_1)$.

The correspondence $M_\omega \rightarrow (M_\omega)^* = M_{\omega^*}$ defines an involutory ring homomorphism

$$* : \text{Qsym}[t_1, t_2, \dots] \rightarrow \text{Qsym}[t_1, t_2, \dots].$$

Remark 3.7. Let $\tau_{\text{Qsym}} : \text{Qsym} \otimes \text{Qsym} \rightarrow \text{Qsym} \otimes \text{Qsym}$ be the ring homomorphism that interchanges the tensor factors $\tau_{\text{Qsym}}(x \otimes y) = y \otimes x$. Then $\Delta(M_\omega^*) = * \otimes * (\tau_{\text{Qsym}} \Delta M_\omega)$.

3.2 Leibnitz-Hopf Algebras

Let R be a commutative associative ring with unity.

Definition 3.8. A *Leibnitz-Hopf algebra* over the ring R is an associative Hopf algebra $\mathcal{H}$ over the ring R with a fixed sequence of a finite or countable number of multiplicative generators H_i , $i = 1, 2, \dots$ satisfying the comultiplication formula

$$\Delta H_n = \sum_{i+j=n} H_i \otimes H_j, \quad H_0 = 1.$$

A *universal Leibnitz-Hopf algebra* $\mathcal{A}$ over the ring R is a Leibnitz-Hopf algebra with the universal property: for any Leibnitz-Hopf algebra $\mathcal{H}$ over the ring R the correspondence $A_i \rightarrow H_i$ defines a Hopf algebra homomorphism.

Consider the free associative Leibnitz-Hopf algebra over the integers $\mathcal{Z} = \mathbb{Z}\langle Z_1, Z_2, \dots \rangle$ in countably many generators Z_i .

Proposition 3.9. $\mathcal{Z} \otimes R$ is a universal Leibnitz-Hopf algebra over the ring R .

Set $\deg Z_i = 2i$. Let us denote by $\mathcal{M}$ the graded dual Hopf algebra over the integers.

It is not difficult to see that $\mathcal{M}$ is precisely the algebra of quasi-symmetric functions over the integers. Indeed, for any composition $\omega = (j_1, \dots, j_k)$ we can define m_ω by the dual basis formula

$$\langle m_\omega, Z_\sigma \rangle = \delta_{\omega, \sigma},$$

where $Z_\sigma = Z_{a_1} \dots Z_{a_l}$ for a composition $\sigma = (a_1, \dots, a_l)$. Then the elements m_ω are multiplied exactly as the quasi-symmetric monomials M_ω .

Let us denote

$$\Phi(t) = 1 + Z_1 t + Z_2 t^2 + \dots = \sum_{k=0}^{\infty} Z_k t^k$$

Then the comultiplication formula is equivalent to

$$\Delta \Phi(t) = \Phi(t) \otimes \Phi(t)$$

Let $\chi : \mathcal{Z} \rightarrow \mathcal{Z}$ be the *antipode*, that is a linear operator, satisfying the property

$$1 \star \chi = \mu \circ (1 \otimes \chi) \circ \Delta = \eta \circ \varepsilon = \mu \circ (\chi \otimes 1) \circ \Delta = \chi \star 1,$$

where $\varepsilon : \mathcal{Z} \rightarrow \mathbb{Z}$, $\varepsilon(1) = 1$, $\varepsilon(Z_\sigma) = 0$, $\sigma \neq \emptyset$, is a *counit*, $\eta : \mathbb{Z} \rightarrow \mathcal{Z}$, $\eta(a) = a \cdot 1$ is a *unite* map, and μ is a multiplication in $\mathcal{Z}$.

Then $\Phi(t)\chi(\Phi(t)) = 1 = \chi(\Phi(t))\Phi(t)$ and $\{\chi(Z_n)\}$ satisfy the recurrent formulas

$$\chi(Z_1) = -Z_1; \quad \chi(Z_{n+1}) + \chi(Z_n)Z_1 + \dots + \chi(Z_1)Z_n + Z_{n+1} = 0, \quad n \geq 1. \quad (6)$$

In fact, since $\Phi(t) = 1 + Z_1 t + \dots$, we can use the previous formula to obtain

$$\begin{aligned} \chi(\Phi(t)) &= \frac{1}{\Phi(t)} = \sum_{i=0}^{\infty} (-1)^i (\Phi(t) - 1)^i; \\ \chi(Z_n) &= \sum_{k=1}^n (-1)^k \sum_{j_1 + \dots + j_k = n} Z_{j_1} \dots Z_{j_k} \quad j_i \geq 1, \quad n \geq 1 \end{aligned}$$

This formula defines χ on generators, and therefore on the whole algebra.

Definition 3.10. Let $\mathcal{Z}^{op}$ be the Hopf algebra opposite to $\mathcal{Z}$, i.e. $\mathcal{Z} = \mathcal{Z}^{op}$ as coalgebras, and the multiplication in $\mathcal{Z}^{op}$ is given by the rule $a \diamond b := b \cdot a$. Then the antiisomorphism

$$\varrho : \mathcal{Z} \rightarrow \mathcal{Z}, \quad Z_{a_1} \dots Z_{a_k} \rightarrow Z_{a_k} \dots Z_{a_1}$$

satisfies the property $\Delta \circ \varrho = (\varrho \otimes \varrho) \circ \Delta$, and defines the Hopf algebra isomorphism $\mathcal{Z} \rightarrow \mathcal{Z}^{op}$.

Proposition 3.11. We have $\varrho^* = *$

Proof. Indeed,

$$\langle \varrho^* M_\omega, Z_\sigma \rangle = \langle M_\omega, \varrho Z_\sigma \rangle = \langle M_\omega, Z_{\sigma^*} \rangle = \delta_{\omega, \sigma^*}.$$

Therefore, $\varrho^* M_\omega = M_{\omega^*} = M_\omega^*$ □

Definition 3.12 ([N]). A (left) *Milnor module* M over the Hopf algebra X is an algebra with unit $1 \in R$ which is also a (left) module over X satisfying

$$x(uv) = \sum x'_n(u)x''_n(v), \quad x \in X, \quad u, v \in M, \quad \Delta x = \sum x'_n \otimes x''_n.$$

Proposition 3.13. The homomorphism $\mathcal{L}_R : \mathcal{Z} \rightarrow \mathcal{D}(R) : Z_k \rightarrow d_k$ defines on the rings $\mathcal{P}$ and $\mathcal{RP}$ the structures of left Milnor modules over the Leibnitz-Hopf algebra $\mathcal{Z}$.

The homomorphism $\mathcal{R}_R : \mathcal{Z}^{op} \rightarrow \mathcal{D}(R) : \mathcal{R}_R = \mathcal{L}_R \circ \varrho$ defines on the rings $\mathcal{P}$ and $\mathcal{RP}$ the structures of right Milnor modules over the Leibnitz-Hopf algebra $\mathcal{Z}$.

Proof. This follows from Proposition 2.49 $\square$

Let us note, that the word $Z_\omega = Z_{j_1} \dots Z_{j_k}$ corresponds under the homomorphism $\mathcal{L}_R$ to the operator $D_\omega = d_{j_1} \dots d_{j_k}$, and under the homomorphism $\mathcal{R}_R$ to the operator $D_{\omega^*} = d_{j_k} \dots d_{j_1}$.

Definition 3.14. Denote by $\mathcal{U}$ the universal Hopf algebra in the category of Leibnitz-Hopf algebras with an antipode $\chi(H_i) = (-1)^i H_i$. It is easy to see that $\mathcal{U} = \mathcal{Z}/J_{\mathcal{U}}$, where the two-sided Hopf ideal $J_{\mathcal{U}}$ is generated by the relations

$$Z_n - Z_1 Z_{n-1} + \dots + (-1)^{n-1} Z_{n-1} Z_1 + (-1)^n Z_n = 0, \quad n \geq 2 \quad (7)$$

These relations can be written in the short form as $\Phi(-t)\Phi(t) = 1$.

It can be shown that $\mathcal{U} \otimes \mathbb{Q} \simeq \mathbb{Q}\langle Z_1, Z_3, Z_5, \dots \rangle$, and $\mathcal{U} \xrightarrow{\text{id} \otimes 1} \mathcal{U} \otimes \mathbb{Q}$ is an embedding.

Corollary 3.15. The mappings $\mathcal{L}_R$ and $\mathcal{R}_R$ define on the rings $\mathcal{P}$ and $\mathcal{RP}$ the structures of left and right Milnor modules over the Leibnitz-Hopf algebra $\mathcal{U}$.

Proof. This follows from Proposition 2.52. $\square$

Since $\varrho(J_{\mathcal{U}}) = J_{\mathcal{U}}$, we see that ϱ induces a correctly defined homomorphism $\varrho: \mathcal{U} \rightarrow \mathcal{U}$, which gives the isomorphism $\mathcal{U} \simeq \mathcal{U}^{op}$.

Since the factor map $\mathcal{Z} \rightarrow \mathcal{Z}/J_{\mathcal{U}}$ is an epimorphism, the dual map $\mathcal{U}^* \rightarrow \mathcal{Z}^* = \mathcal{Qsym}$ is an embedding. Then the subring $\mathcal{U}^* \subset \mathcal{Qsym}$ is invariant under the involution $\varrho^* = *$.

Definition 3.16. A universal commutative Leibnitz-Hopf algebra $\mathcal{C} = \mathbb{Z}[C_1, C_2, \dots]$ is a free commutative polynomial Leibnitz-Hopf algebra in generators C_i of degree $2i$. We have $\mathcal{C} = \mathcal{Z}/J_{\mathcal{C}}$, where the ideal $J_{\mathcal{C}}$ is generated by the relations $Z_i Z_j - Z_j Z_i$.

It is a self-dual Hopf algebra and the graded dual Hopf algebra is naturally isomorphic to the algebra of symmetric functions $\mathbb{Z}[\sigma_1, \sigma_2, \dots] = \text{Sym}[t_1, t_2, \dots] \subset \mathcal{Qsym}[t_1, t_2, \dots]$ generated by the symmetric monomials

$$\sigma_i = M_{\omega_i} = \sum_{l_1 < \dots < l_i} t_{l_1} \dots t_{l_i},$$

where $\omega_i = \underbrace{(1, \dots, 1)}_i$.

The isomorphism $\mathcal{C} \simeq \mathcal{C}^*$ is given by the correspondence $C_i \rightarrow \sigma_i$.

Definition 3.17. Denote by $\mathcal{US}$ the universal Hopf algebra in the category of commutative Leibnitz-Hopf algebras with an antipode $\chi(H_i) = (-1)^i H_i$. It is easy to see that $\mathcal{US} = \mathcal{Z}/J_{\mathcal{US}}$, where the two-sided ideal $J_{\mathcal{US}}$ is generated by the relations (7), and the commutators $Z_i Z_j - Z_j Z_i$. Then the relations (7) for odd n are trivial, while for even n we have

$$2Z_{2k} = 2 \sum_{i=1}^{k-1} (-1)^{i-1} Z_i Z_{2k-i} + (-1)^{k-1} Z_k^2$$

It can be shown that $\mathcal{US} \otimes \mathbb{Q} \simeq \mathbb{Q}[Z_1, Z_3, Z_5, \dots]$.

3.3 Lie-Hopf Algebras

Definition 3.18. A Lie-Hopf algebra over the ring R is an associative Hopf algebra $\mathfrak{L}$ with a fixed sequence of a finite or countable number of multiplicative generators L_i , $i = 1, 2, \dots$ satisfying the comultiplication formula

$$\Delta L_i = 1 \otimes L_i + L_i \otimes 1, \quad L_0 = 1.$$

A universal Lie-Hopf algebra over the ring R is a Lie-Hopf algebra $\mathcal{A}$ satisfying the universal property: for any Lie-Hopf algebra $\mathfrak{L}$ over the ring R the correspondence $A_i \rightarrow L_i$ defines a Hopf algebra homomorphism $\mathcal{A} \rightarrow \mathfrak{L}$.

Consider the free associative Lie-Hopf algebra $\mathcal{W}$ over the integers $\mathbb{Z}\langle W_1, W_2, \dots \rangle$ in countably many variables W_i .

Proposition 3.19. $\mathcal{W} \otimes R$ is a universal Lie-Hopf algebra over the ring R .

Let us set $\deg W_i = 2i$. Then the graded dual Hopf algebra is denoted by $\mathcal{N}$. This is the so-called *shuffle algebra*.

The antipode χ in $\mathcal{W}$ has a very simple form $\chi(W_i) = -W_i$.

3.4 Lyndon Words

A well-known theorem in the theory of free Lie algebras (see, for example, [Re]) states that the algebra $\mathcal{N} \otimes \mathbb{Q}$ is a commutative free polynomial algebra in the so-called *Lyndon words*.

Definition 3.20. Let us denote by $[a_1, \dots, a_n]$ an element of $\mathbb{N}^*$, that is the word over $\mathbb{N}$, consisting of symbols $a_1, \dots, a_n$, $a_i \in \mathbb{N}$. Let us order the words from $\mathbb{N}^*$ lexicographically, where any symbol is larger than nothing, that is $[a_1, \dots, a_n] > [b_1, \dots, b_m]$ if and only if there is an i such that $a_1 = b_1, \dots, a_{i-1} = b_{i-1}, a_i > b_i, 1 \leq i \leq \min\{m, n\}$, or $n > m$ and $a_1 = b_1, \dots, a_m = b_m$.

A *proper tail* of a word $[a_1, \dots, a_n]$ is a word of the form $[a_i, \dots, a_n]$ with $1 < i \leq n$. (The empty word and one-symbol word have no proper tails.)

A word is *Lyndon* if all its proper tails are larger than the word itself. For example, the words $[1, 1, 2]$, $[1, 2, 1, 2, 2]$, $[1, 3, 1, 5]$ are Lyndon and the words $[1, 1, 1, 1]$, $[1, 2, 1, 2]$, $[2, 1]$ are not Lyndon. The set of Lyndon words is denoted by LYN .

The same definitions make sense for any totally ordered set, for example, for the set $\{1, 2\}$ or for the set of all odd positive integers.

The role of Lyndon words is described by the following theorem:

Theorem (Chen-Fox-Lyndon Factorization [CFL]). *Every word w in $\mathbb{N}^*$ factors uniquely into a decreasing concatenation product of Lyndon words*

$$w = u_1 * u_2 * \dots * u_k, \quad u_i \in \text{LYN}, \quad u_1 \geq u_2 \geq \dots \geq u_k$$

For example, $[1, 1, 1, 1] = [1] * [1] * [1] * [1]$, $[1, 2, 1, 2] = [1, 2] * [1, 2]$, $[2, 1] = [2] * [1]$.

The algebra $\mathcal{N}$ is additively generated by the words in $\mathbb{N}^*$. The word $w = [a_1, \dots, a_n]$ corresponds to the function

$$\langle [a_1, \dots, a_n], W_\sigma \rangle = \delta_{w, \sigma}$$

where σ is a composition $\sigma = (b_1, \dots, b_l)$, $W_\sigma = W_{b_1} \dots W_{b_l}$, and for the word $[a_1, \dots, a_n]$

$$\delta_{w, \sigma} = \begin{cases} 1, & \sigma = (a_1, \dots, a_n); \\ 0, & \text{else} \end{cases}$$

The multiplication in the algebra $\mathcal{N}$ is the so-called *shuffle multiplication*:

$$[a_1, \dots, a_n] \times_{sh} [a_{n+1}, \dots, a_{m+n}] = \sum_{\sigma} [a_{\sigma(1)}, \dots, a_{\sigma(n)}, a_{\sigma(n+1)}, \dots, a_{\sigma(n+m)}],$$

where σ runs over all the substitutions $\sigma \in S_{m+n}$ such that

$$\sigma^{-1}(1) < \dots < \sigma^{-1}(n) \text{ and } \sigma^{-1}(n+1) < \dots < \sigma^{-1}(n+m).$$

For example,

$$\begin{aligned} [1] \times_{sh} [1] &= [1, 1] + [1, 1] = 2[1, 1]; \\ [1] \times_{sh} [2, 3] &= [1, 2, 3] + [2, 1, 3] + [2, 3, 1]; \\ [1, 2] \times_{sh} [1, 2] &= [1, 2, 1, 2] + [1, 1, 2, 2] + [1, 1, 2, 2] + [1, 1, 2, 2] + [1, 1, 2, 2] + [1, 2, 1, 2] = 2[1, 2, 1, 2] + 4[1, 1, 2, 2]. \end{aligned}$$

There is a well-known shuffle algebra structure theorem:

Theorem. $\mathcal{N} \otimes \mathbb{Q} = \mathbb{Q}[\text{LYN}]$, the free commutative algebra over $\mathbb{Q}$ in the symbols from LYN .

The proof follows from the following theorem concerning shuffle products in connection with Chen-Fox-Lyndon factorization.

Theorem. *Let w be a word on the natural numbers and let $w = u_1 * u_2 * \cdots * u_m$ be its Chen-Fox-Lyndon factorization. Then all words that occur with nonzero coefficient in the shuffle product $u_1 \times_{sh} u_2 \times_{sh} \cdots \times_{sh} u_m$ are lexicographically smaller or equal to w , and w occurs with a nonzero coefficient in this product.*

Given this result it is easy to prove the shuffle algebra structure theorem in the case of arbitrary totally ordered subset $M = \{m_1, m_2, \dots\} \subset \mathbb{N}$. Let us denote by $\mathcal{W}_M$ the free associative Lie-Hopf algebra $\mathbb{Z}\langle W_{m_1}, W_{m_2}, \dots \rangle$, let $\mathcal{N}_M$ be its graded dual algebra, and LYN_M be the corresponding set of all Lyndon words. We need to prove that $\mathcal{N}_M \otimes \mathbb{Q} = \mathbb{Q}[\text{LYN}_M]$.

Let m_1 be the minimal number in M . The smallest word $[m_1]$ is Lyndon.

Given a word w we can assume by induction that all the words lexicographically smaller than w have been written as polynomials in the elements of LYN_M . Take the Chen-Fox-Lyndon factorization $w = u_1 * u_2 * \cdots * u_m$ of w and consider, using the preceding theorem,

$$u_1 \times_{sh} u_2 \times_{sh} \cdots \times_{sh} u_m = aw + (\text{remainder}).$$

By the theorem the coefficient a is nonzero and all the words in (remainder) are lexicographically smaller than w , hence they belong to $\mathbb{Q}[\text{LYN}_M]$. Therefore $w \in \mathbb{Q}[\text{LYN}_M]$. This proves generation. Since each monomial in $(2n)$ -th graded component of $\mathcal{N} \otimes \mathbb{Q}$ has a unique Chen-Fox-Lyndon decomposition, the number of monomials in $\mathbb{Q}[\text{LYN}_M]$ of graduation $2n$ is equal to the number of monomials of the same graduation in $\mathcal{N}_M \otimes \mathbb{Q}$, monomials in Lyndon words are linearly independent. This proves that Lyndon words are algebraically independent.

Corollary 3.21. *The graded dual algebras to the free associative Lie-Hopf algebras $\mathcal{W}_{12} = \mathbb{Z}\langle W_1, W_2 \rangle$ and $\mathcal{W}_{\text{odd}} = \mathbb{Z}\langle W_1, W_3, W_5, \dots \rangle$ are free polynomial algebras in Lyndon words LYN_{12} and LYN_{odd} respectively.*

The correspondence

$$1 + Z_1 t + Z_2 t^2 + \cdots = \exp(W_1 t + W_2 t^2 + \dots);$$

defines an isomorphism of Hopf algebras $\mathcal{Z} \otimes \mathbb{Q} \simeq \mathcal{W} \otimes \mathbb{Q}$.

However it is not true that $\mathcal{N}$ is a free polynomial commutative algebra over the integers.

4 Topological realization of Hopf algebras

In [BR] A. Baker and B. Richter showed that the ring of quasi-symmetric functions has a very nice topological interpretation.

We will consider CW -complexes X and their homology $H_*(X)$ and cohomology $H^*(X)$ with integer coefficients.

Let us assume that homology groups $H_*(X)$ have no torsion.

The diagonal map $X \rightarrow X \times X$ defines in $H_*(X)$ the structure of a graded coalgebra with the comultiplication $\Delta: H_*(X) \rightarrow H_*(X) \otimes H_*(X)$ and the dual structure of a graded algebra in $H^*(X)$ with the multiplication $\Delta^*: H^*(X) \otimes H^*(X) \rightarrow H^*(X)$.

In the case when X is an H -space with the multiplication $\mu: X \times X \rightarrow X$ we obtain the Pontryagin product $\mu_*: H_*(X) \otimes H_*(X) \rightarrow H_*(X)$ in the coalgebra $H_*(X)$ and the corresponding structure of a graded Hopf algebra on $H_*(X)$. The cohomology ring $H^*(X)$ obtains the structure of a graded dual Hopf algebra with the diagonal mapping $\mu^*: H^*(X) \rightarrow H^*(X) \otimes H^*(X)$.

For any space Y the loop space $X = \Omega Y$ is an H -space. A continuous mapping $f: Y_1 \rightarrow Y_2$ induces a mapping of H -spaces $\Omega f: X_1 \rightarrow X_2$, where $X_i = \Omega Y_i$. Thus for any space Y such that $X = \Omega Y$ has no torsion in integral homology we obtain the Hopf algebra $H_*(X)$. This correspondence is functorial, that is any continuous mapping $f: Y_1 \rightarrow Y_2$ induces a Hopf algebra homomorphism $f_*: H_*(X_1) \rightarrow H_*(X_2)$.

By the Bott-Samelson theorem [BS], $H_*(\Omega \Sigma X)$ is the free associative algebra $T(\tilde{H}_*(X))$ generated by $\tilde{H}_*(X)$. This construction is functorial, that is a continuous mapping $f: X_1 \rightarrow X_2$ induces a ring homomorphism of the corresponding tensor algebras arising from the mapping $f_*: \tilde{H}_*(X_1) \rightarrow \tilde{H}_*(X_2)$. Denote

elements of $H_*(\Omega\Sigma X)$ by $(a_1|\dots|a_n)$, where $a_i \in \tilde{H}_*(X)$. Since the diagonal mapping $\Sigma X \rightarrow \Sigma X \times \Sigma X$ gives the H -map $\Delta: \Omega\Sigma X \rightarrow \Omega\Sigma X \times \Omega\Sigma X$, it follows from the Eilenberg-Zilber theorem that

$$\Delta_*(a_1|\dots|a_n) = (\Delta_*a_1|\dots|\Delta_*a_n)$$

where $(a_1 \otimes b_1|a_2 \otimes b_2) = (a_1|a_2) \otimes (b_1|b_2)$.

There is a nice combinatorial model for any topological space of the form $\Omega\Sigma X$ with X connected, namely the James construction JX on X . After one suspension this gives rise to a splitting

$$\Sigma\Omega\Sigma X \sim \Sigma JX \sim \bigvee_{n \geq 1} \Sigma X^{(n)},$$

where $X^{(n)}$ denotes the n -fold smash power of X .

Example 4.1. There exists a homotopy equivalence

$$\Sigma\Omega\Sigma S^2 \rightarrow \bigvee_{n \geq 1} \Sigma(S^2)^{(n)} \simeq \Sigma\left(\bigvee_{n \geq 1} S^{2n}\right).$$

Therefore there exists an H -map $\Omega\Sigma(\Omega\Sigma S^2) \rightarrow \Omega\Sigma\left(\bigvee_{n \geq 1} S^{2n}\right)$ that is a homotopy equivalence.

Using these classical topological results let us describe topological realizations of the Hopf algebras we study in this work.

We have the following Hopf algebras:

- I.** 1. $H_*(\mathbb{C}P^\infty)$ is a divided power algebra $\mathbb{Z}[u_1, u_2, \dots]/I$, where the ideal I is generated by the relations $u_i u_j - \binom{i+j}{i} u_{i+j}$, with the comultiplication

$$\Delta u_n = \sum_{k=0}^n u_k \otimes u_{n-k}; \quad (8)$$

2. $H^*(\mathbb{C}P^\infty) = \mathbb{Z}[u]$ – a polynomial ring with the comultiplication

$$\Delta u = 1 \otimes u + u \otimes 1;$$

- II.** 1. $H_*(\Omega\Sigma\mathbb{C}P^\infty) \simeq T(\tilde{H}_*(\mathbb{C}P^\infty)) = \mathbb{Z}\langle u_1, u_2, \dots \rangle$ with u_i being non-commuting variables of degree $2i$. Thus there is an isomorphism of rings

$$H_*(\Omega\Sigma\mathbb{C}P^\infty) \simeq \mathcal{Z}$$

under which u_n corresponds to Z_n .

The coproduct Δ on $H_*(\Omega\Sigma\mathbb{C}P^\infty)$ induced by the diagonal in $\Omega\Sigma\mathbb{C}P^\infty$ is compatible with the one in $\mathcal{Z}$:

$$\Delta u_n = \sum_{i+j=n} u_i \otimes u_j$$

So there is an isomorphism of graded Hopf algebras. This gives a geometric interpretation for the antipode χ in $\mathcal{Z}$: in $H_*(\Omega\Sigma\mathbb{C}P^\infty)$ it arises from the time-inversion of loops.

2. $H^*(\Omega\Sigma\mathbb{C}P^\infty)$ is the graded dual Hopf algebra to $H_*(\Omega\Sigma\mathbb{C}P^\infty)$.

Theorem. *[BR] There is an isomorphism of graded Hopf algebras:*

$$H_*(\Omega\Sigma\mathbb{C}P^\infty) \simeq \mathcal{Z} = \mathbb{Z}\langle Z_1, Z_2, \dots \rangle, \quad H^*(\Omega\Sigma\mathbb{C}P^\infty) \simeq \mathcal{M} = \text{Qsym}[t_1, t_2, \dots].$$

- III.** $H_*(BU) \simeq H^*(BU) \simeq \mathbb{Z}[\sigma_1, \sigma_2, \dots] \simeq \mathcal{C}$. It is a self-dual Hopf algebra of symmetric functions. In the cohomology σ_i are represented by Chern classes.

IV. 1. $H_*(\Omega\Sigma S^2) = H_*(\Omega S^3) = \mathbb{Z}[w]$ – a polynomial ring with $\deg w = 2$ and the comultiplication

$$\Delta w = 1 \otimes w + w \otimes 1;$$

2. $H^*(\Omega\Sigma S^2) = \mathbb{Z}[u_n]/I$ is a divided power algebra. Thus $H^*(\Omega\Sigma S^2) \simeq H_*(\mathbb{C}P^\infty)$.

V. $H_*(\Omega\Sigma(\Omega\Sigma S^2)) \simeq \mathbb{Z}\langle w_1, w_2, \dots \rangle$. It is a free associative Hopf algebra with the comultiplication

$$\Delta w_n = \sum_{k=0}^n \binom{n}{k} w_k \otimes w_{n-k} \quad (9)$$

VI. $H_*\left(\Omega\Sigma\left(\bigvee_{n \geq 1} S^{2n}\right)\right) \simeq \mathbb{Z}\langle \xi_1, \xi_2, \dots \rangle$. It is a free associative algebra and has the structure of a graded Hopf algebra with the comultiplication

$$\Delta \xi_n = 1 \otimes \xi_n + \xi_n \otimes 1. \quad (10)$$

Therefore, $H_*\left(\Omega\Sigma\left(\bigvee_{n \geq 1} S^{2n}\right)\right)$ gives a topological realization of the universal Lie-Hopf algebra $\mathcal{W}$.

The homotopy equivalence

$$a: \Omega\Sigma\left(\bigvee_{n \geq 1} S^{2n}\right) \longrightarrow \Omega\Sigma(\Omega\Sigma S^2)$$

induces an isomorphism of graded Hopf algebras

$$a_*: \mathbb{Z}\langle \xi_1, \xi_2, \dots \rangle \longrightarrow \mathbb{Z}\langle w_1, w_2, \dots \rangle,$$

and its algebraic form is determined by the conditions:

$$\Delta a_* \xi_n = (a_* \otimes a_*)(\Delta \xi_n).$$

For example, $a_* \xi_1 = w_1$, $a_* \xi_2 = w_2 - w_1|w_1$, $a_* \xi_3 = w_3 - 3w_2|w_1 + 2w_1|w_1|w_1$.

Thus using topological results we have obtained that two Hopf algebra structures on the free associative algebra with the comultiplications (9) and (10) are isomorphic over $\mathbb{Z}$.

This result is interesting from the topological point of view, since the elements $(w_n - a_* \xi_n)$ for $n \geq 2$ are obstructions to the desuspension of the homotopy equivalence

$$\Sigma(\Omega\Sigma S^2) \rightarrow \Sigma\left(\bigvee_{n \geq 1} S^{2n}\right).$$

We have the commutative diagram:

$$\begin{array}{ccccc} \Omega\Sigma(\Omega\Sigma S^2) & \xrightarrow{\Omega\Sigma k} & \Omega\Sigma\mathbb{C}P^\infty & & \\ \uparrow & & \uparrow & \searrow j & \\ \Omega\Sigma S^2 & \xrightarrow{k} & \mathbb{C}P^\infty & \xrightarrow{i} & BU \xrightarrow{\det} \mathbb{C}P^\infty \end{array}$$

Here

- i is the inclusion $\mathbb{C}P^\infty = BU(1) \subset BU$,
- j arises from the universal property of $\Omega\Sigma\mathbb{C}P^\infty$ as a free H -space;
- the mapping $\mathbb{C}P^\infty \rightarrow \Omega\Sigma\mathbb{C}P^\infty$ is induced by the identity map $\Sigma\mathbb{C}P^\infty \rightarrow \Sigma\mathbb{C}P^\infty$,
- the mapping $k: \Omega\Sigma S^2 \rightarrow \mathbb{C}P^\infty$ corresponds to the generator of $H^2(\Omega\Sigma S^2) = \mathbb{Z}$.
- $\det$ is the mapping of the classifying spaces $BU \rightarrow BU(1)$ induced by the mappings $\det: U(n) \rightarrow U(1)$, $n = 1, 2, \dots$.

As we have mentioned, $H^*(BU)$ as a Hopf algebra is isomorphic to the algebra of symmetric functions (in the generators – Chern classes) and is self-dual.

$$H^*(BU) \simeq H_*(BU) \simeq \text{Sym}[t_1, t_2, \dots] = \mathbb{Z}[\sigma_1, \sigma_2, \dots].$$

Since $\mathbb{C}P^\infty$ gives rise to the algebra generators in $H_*(BU)$, j_* is an epimorphism on homology, and j^* is a monomorphism on cohomology.

- j_* corresponds to the factorization of the non-commutative polynomial algebra $\mathcal{Z}$ by the commutation relations $Z_i Z_j - Z_j Z_i$ and sends Z_i to σ_i .
- j^* corresponds to the inclusion $\text{Sym}[t_1, t_2, \dots] \subset \text{Qsym}[t_1, t_2, \dots]$ and sends σ_i to M_{ω_i} , where $\omega_i = \underbrace{(1, 1, \dots, 1)}_i$.
- k^* sends u to u_1 and defines an embedding of the polynomial ring $\mathbb{Z}[u]$ into the divided power algebra $\mathbb{Z}[u_n]/I$.
- the composition $\det_* \circ j_*$ maps the algebra $\mathcal{Z}$ to the divided power algebra $H_*(\mathbb{C}P^\infty)$. This corresponds to the mapping $Z_i \rightarrow d_k|_{\mathcal{P}_s}$ of $\mathcal{Z}$ to $\mathcal{D}(\mathcal{P}_s)$.

5 Structure of $\mathcal{D}$

Theorem 5.1. *The homomorphism $\mathcal{L}_R : \mathcal{Z} \rightarrow \mathcal{D}(R) : Z_k \rightarrow d_k$ induces a ring isomorphism:*

$$\mathcal{D}(R) \simeq \mathcal{U} = \mathcal{Z}/J_{\mathcal{U}},$$

where $\mathcal{U}$ is a universal Leibnitz-Hopf algebra with the antipode $\chi(U_i) = (-1)^i U_i$.

Proof. Let us prove the following lemma:

Lemma 5.2. *Let $D \in \mathcal{D}(R)$ be an operator of graduation $2k$. Then for each space R^{2n} , $n \geq k$ there is a unique representation*

$$D = u(d_2, d_3, \dots, d_k) + dw(d_2, d_3, \dots, d_{k-1}), \quad (11)$$

and there is a unique representation

$$D = u'(d_2, d_3, \dots, d_k) + w'(d_2, \dots, d_{k-1})d, \quad (12)$$

where u, u', w, w' are polynomials.

Proof. For $k = 0$ and 1 it is evident.

Since $2d_2 = d^2$, this is true for $k = 2$. Let $k \geq 3$.

Using the relations (2.52) we obtain

$$dd_i = (-1)^i d_i d + (d_2 d_{i-1} - d_3 d_{i-2} + \dots + (-1)^{i-1} d_{i-1} d_2) + (1 + (-1)^{i+1}) d_{i+1}$$

So the expressions (11) and (12) exist.

Let $u(d_2, d_3, \dots, d_k) + dw(d_2, \dots, d_{k-1}) = 0$, where $u = \sum_{|\omega|=k} a_\omega D_\omega$, and $w = \sum_{|\omega|=k-1} b_\omega D_\omega$.

Then for any n -dimensional polytope P^n , $n \geq k$ we have

$$\xi_1(u + dw)P^n = 0, \quad \varepsilon_1(u + dw)P^n = 0$$

Since $u + dw$ has degree $2k$, this equality can be written as

$$\sum_{|\omega|=k} a_\omega f_{n-k, n-k+j_1, n-k+j_1+j_2, \dots, n-j_l}(P^n) + \sum_{|\omega|=k-1} b_\omega f_{n-k, n-k+1, n-k+1+j_1, \dots, n-j_l}(P^n). \quad (13)$$

Using the generalized Dehn-Sommerville equations we obtain

$$f_{n-k, n-k+1, n-k+1+j_1, \dots, n-j_l} = (-1)^{n-k-1} \left(\sum_{j=0}^{n-k-1} (-1)^j f_{j, n-k+1, n-k+1+j_1, \dots, n-j_l} \right) + (1 + (-1)^{n-k}) f_{n-k+1, n-k+1+j_1, \dots, n-j_l}$$

All the sets $\{n-k, n-k+j_1, n-k+j_1+j_2, \dots, n-j_l\}$, $\{j, n-k+1, n-k+j_1, \dots, n-j_l\}$ and $\{n-k+1, n-k+1+j_1, \dots, n-j_l\}$ for all ω are different and belong to Ψ^n .

In the case of $\mathcal{RP}$ and P^{k-1} we obtain that $n = k-1$, therefore $n-k = -1$ in the expression (13) vanishes and we obtain

$$\sum_{|\omega|=k} a_\omega f_{j_1-1, j_1+j_2-1, \dots, n-j_l}(P^n) + \sum_{|\omega|=k-1} b_\omega f_{0, j_1, \dots, n-j_l}(P^n).$$

Again all the sets $\{j_1-1, j_1+j_2-1, \dots, n-j_l\}$, $\{0, j_1, \dots, n-j_l\}$ for all ω are different and belong to Ψ^n .

Since the vectors $\{f_S(Q), S \in \Psi^n\}$, $Q \in \Omega^n$ are linearly independent, we obtain that all a_ω and b_ω are equal to 0, so the representation (11) is unique.

We obtain that the monomials $D_\omega = d_{j_1} \dots d_{j_l}$, $|\omega| = k$, $j_i \geq 2$ and $dD_\omega = dd_{j_1} \dots d_{j_l}$ $|\omega| = k-1$, $j_i \geq 2$ form a basis of the abelian group $\mathcal{D}(\mathbb{R}^{2n}, \mathbb{R}^{2(n-k)})$.

But each monomial dD_ω , $|\omega| = k$, $j_i \geq 2$ can be expressed as an integer combination of the monomials $D_{\omega'}$ and $D_{\omega'}d$. So the monomials D_ω and $D_\omega d$ also form a basis. This proves the second part of the lemma. $\square$

Now let us prove the theorem. The mapping $\mathcal{L}_R : \mathcal{Z} \rightarrow \mathcal{D}(\mathbb{R}) : Z_i \rightarrow d_i$ is an epimorphism. Let $z \in \mathcal{Z}$ such that $\deg z = 2k$ and $\mathcal{L}_R z = 0$.

$$z = \sum_{|\omega|=k} a_\omega Z_\omega.$$

We know that

$$Z_1 Z_{k-1} = (-1)^{k-1} Z_{k-1} Z_1 + \sum_{i=2}^{k-2} (-1)^i Z_i Z_{k-i} + (1 + (-1)^k) Z_k - \sum_{i=0}^k (-1)^i Z_i Z_{k-i}.$$

So

$$z = \sum_{|\omega|=k, j_i \geq 2} a'_\omega Z_\omega + \sum_{|\omega|=k-1, j_i \geq 2} b_\omega Z_\omega Z_1 + z',$$

where $z' \in J_{\mathcal{U}}$. Since $\mathcal{L}_R z = \mathcal{L}_R z' = 0$, we obtain

$$\sum_{|\omega|=k, j_i \geq 2} a'_\omega D_\omega + \sum_{|\omega|=k-1, j_i \geq 2} b_\omega D_\omega d = 0,$$

Lemma 5.2 implies that all the coefficients a'_ω and b_ω are equal to 0, so $z = z' \in J_{\mathcal{U}}$. This proves that $\mathcal{D}(\mathbb{R}) \simeq \mathcal{Z}/J_{\mathcal{U}}$. $\square$

Remark 5.3. Now we see that the correspondence $d_k \rightarrow d_k$ defines a natural isomorphism between $\mathcal{D}(\mathcal{RP})$ and $\mathcal{D}(\mathcal{P})$. Therefore from this moment we will often use the notation $\mathcal{D}$ for both these algebras, and the notations $\mathcal{L}$, $\mathcal{R}$ for the homomorphisms $\mathcal{L}_R$ and $\mathcal{R}_R$.

Corollary 5.4. *The comultiplication $d_k \rightarrow \sum_{i+j=k} d_i \otimes d_j$ defines on $\mathcal{D}$ the structure of a Leibnitz-Hopf algebra over the ring $\mathbb{Z}$. This Hopf algebra $\mathcal{D}$ is isomorphic to the Hopf algebra $\mathcal{U}$, universal in the category of Leibnitz-Hopf algebras with the antipode $\chi(U_i) = (-1)^i U_i$.*

Both $\mathcal{P}$ and $\mathcal{RP}$ have natural left Milnor module structures over $\mathcal{D}$

It is evident that $D_\omega|_{\mathbb{R}^{[2n]}} = 0$ for $|\omega| > n$. The following corollary says that the other relations between operators in $\mathcal{D}$ on the abelian group $\mathbb{R}^{[2n]}$ are the same as on the whole ring $\mathbb{R}$.

Corollary 5.5. *We have*

$$\mathcal{D}(\mathbb{R}^{[2n]}) = \mathcal{D}(\mathbb{R})/J_n$$

where the ideal J_n is generated by all the operators D_ω , $|\omega| > n$.

Proof. As in the proof of Theorem 5.1 using Lemma 5.2 it is easy to show that $\mathcal{D}(\mathbb{R}^{[2n]}) \simeq \mathcal{Z}/J_{\mathcal{D}(\mathbb{R}^{[2n]})}$, where the ideal $J_{\mathcal{D}(\mathbb{R}^{[2n]})}$ is generated by the relations $\Phi(-t)\Phi(t) = 1$ and $Z_\omega = 0$ for $|\omega| > n$. $\square$

Corollary 5.6. *The operators $d_2, d_3, d_4, \dots$ are algebraically independent.*

Corollary 5.7. *Rank of the $(2n)$ -th graded component of the ring $\mathcal{D}$ is equal to the $(n-1)$ -th Fibonacci number c_{n-1} .*

Proof. At first, let us calculate rank rk_n of the $(2n)$ -th graded component of the ring $\mathbb{Z}\langle d_2, d_3, d_4, \dots \rangle$.

$$\text{rk}_0 = 1, \text{rk}_1 = 0, \text{rk}_2 = 1, \text{rk}_3 = 1, \text{rk}_4 = 2, \text{rk}_5 = 3, \dots$$

It is easy to see that there is a recursive relation

$$\text{rk}_{n+1} = \text{rk}_{n-1} + \text{rk}_{n-2} + \dots + \text{rk}_2 + 1, \quad n \geq 3$$

Then $\text{rk}_{n+1} = \text{rk}_{n-1} + \text{rk}_n$. Since $\text{rk}_2 = \text{rk}_3 = 1$, we obtain $\text{rk}_n = c_{n-2}$, $n \geq 2$. At last, rank of the $(2n)$ -th graded component of the ring $\mathcal{D}$ is equal to $\text{rk}_{n-1} + \text{rk}_n = c_{n-3} + c_{n-2} = c_{n-1}$ for $n \geq 3$. It is easy to see that for $n = 1$ and 2 it is also true. $\square$

Let us remind that on the ring of simple polytopes $d_k|_{\mathcal{P}_s} = \frac{d^k}{k!}|_{\mathcal{P}_s}$, so the ring $\mathcal{D}(\mathcal{P}_s)$ is isomorphic to the divided power ring $d_k d_l = \binom{k+l}{k} d_{k+l}$.

Proposition 5.8.

$$\mathcal{D} \otimes \mathbb{Q} = \mathbb{Q}\langle d_1, d_3, d_5, \dots \rangle,$$

The inclusion $\mathcal{D} \subset \mathcal{D} \otimes \mathbb{Q}$ is an embedding, and the operators d_{2k} are expressed in terms of the operators $d_1, d_3, \dots, d_{2k-1}$ by the formulas

$$d_{2k} = \sum_{i=1}^k (-1)^{i-1} \frac{\binom{2i-2}{i-1}}{i 2^{2i-1}} \left(\sum_{j_1+j_2+\dots+j_{2i}=i+k, j_i \geq 1} d_{2j_1-1} \dots d_{2j_{2i}-1} \right). \quad (14)$$

Proof. Let us denote

$$a(t) = \sum_{k \geq 0} d_{2k} t^{2k}, \quad b(t) = \sum_{k \geq 0} d_{2k+1} t^{2k+1}.$$

Then

$$\begin{aligned} \Phi(t) &= a(t) + b(t) & \Phi(-t) &= a(t) - b(t); \\ a(t)b(t) &= \frac{\Phi(t) + \Phi(-t)}{2} \cdot \frac{\Phi(t) - \Phi(-t)}{2} = \frac{\Phi(t)^2 - \Phi(-t)^2}{4} = b(t)a(t). \end{aligned}$$

Then the relation $\Phi(-t)\Phi(t) = 1$ is equivalent to the relations

$$\begin{aligned} a(t)^2 - b(t)^2 &= 1; \\ a(t)b(t) &= b(t)a(t). \end{aligned}$$

Therefore $a(t) = \sqrt{1+b(t)^2}$ and the formula (14) is valid. Consequently all the operators d_{2k} are expressed as polynomials in $d_1, d_3, \dots$ with rational coefficients. For example,

$$d_2 = \frac{d^2}{2}, \quad d_4 = \frac{dd_3 + d_3d}{2} - \frac{d^4}{8}.$$

This means that the algebra $\mathcal{D} \otimes \mathbb{Q}$ is generated by $d_1, d_3, d_5, \dots$. On the other hand, let us calculate the number of the monomials $d_{2j_1-1} \dots d_{2j_k-1}$ with $(2j_1-1) + \dots + (2j_k-1) = n$. Denote this number by l_n . Then

$$l_0 = 1, \quad l_1 = 1, \quad l_2 = 1, \quad l_3 = 2, \quad l_4 = 3, \dots,$$

and there is a recursive formula $l_{n+1} = l_n + l_{n-2} + l_{n-4} + \dots$ for $n \geq 2$. Hence $l_{n+1} = l_n + l_{n-1}$ for $n \geq 2$, and $l_n = c_{n-1}$ for $n \geq 1$.

We see that the number of monomials of degree $2n$ is equal to dimension of the $(2n)$ -th graded component according to Corollary 5.7. This implies that they are linearly independent over the rationals. Therefore $d_1, d_3, d_5, \dots$ are algebraically independent.

At last, the inclusion $\mathcal{D} \subset \mathcal{D} \otimes \mathbb{Q}$ is an embedding, since $\mathcal{D}$ is torsion-free. $\square$

Definition 5.9. Let us define operators s_k by the formula

$$s(t) = s_1 t + s_2 t^2 + s_3 t^3 + \dots = \log \Phi(t).$$

Then $s_1 = d_1$, $s_2 = 0$, $s_3 = d_3 - \frac{d^3}{6}$, and so on.

The relation $\Phi(-t)\Phi(t) = 1$ turns into the relation $s(-t) + s(t) = 0$. Thus $s_{2k} = 0$ for all k .

Also we have $\Delta s(t) = 1 \otimes s(t) + s(t) \otimes 1$, so each operator s_{2k-1} is a derivation.

Proposition 5.10. *There is an isomorphism of Hopf algebras*

$$\mathcal{D} \otimes \mathbb{Q} = \mathbb{Q}\langle s_1, s_3, s_5, \dots \rangle,$$

where $\mathbb{Q}\langle s_1, s_3, s_5, \dots \rangle$ is a free Lie-Hopf algebra in the generators of degree $\deg s_{2k-1} = 2(2k-1)$, $k \geq 1$, the comultiplication $\Delta s_{2k-1} = 1 \otimes s_{2k-1} + s_{2k-1} \otimes 1$, and the antipode $\chi(s_{2k-1}) = -s_{2k-1}$

6 Generalized f -polynomial

6.1 Definition

The mapping $\Phi : \mathcal{P} \rightarrow \mathcal{P}[t]$ is a ring homomorphism. If we set $\Phi(t_2)t_1 = t_1$, then the operator $\Phi(t_2)$ defines a ring homomorphism $\mathcal{P} \rightarrow \mathcal{P}[t_1, t_2] : P \rightarrow \Phi(t_2)\Phi(t_1)P$,

Thus for each $n \geq 0$ we can define the ring homomorphism $\Phi_n : \mathcal{P} \rightarrow \mathcal{P}[[t_1, \dots, t_n]]$ as a composition:

$$\Phi_n(t_1, \dots, t_n)P = \Phi(t_n) \dots \Phi(t_1)P.$$

Definition 6.1. Let P^n be an n -dimensional polytope. Let us define a quasi-symmetric function

$$f_r(\alpha, t_1, \dots, t_r)(P^n) = \xi_\alpha \Phi(t_r) \dots \Phi(t_1)P^n = \alpha^n + \sum_{k=1}^{\min\{r, n\}} \sum_{0 \leq a_1 < \dots < a_k \leq n-1} f_{a_1, \dots, a_k} \alpha^{a_1} M_{(n-a_k, \dots, a_2-a_1)}.$$

For $n = 1$ we obtain $f_1(\alpha, t_1)(P^n) = \alpha^n + f_{n-1} \alpha^{n-1} t_1 + \dots + f_0 t_1^n$ is a homogeneous f -polynomial in two variables ([Buch]). So the polynomial f_n is a generalization of the f -polynomial.

Consider the increasing sequence of rings

$$\mathbb{Z}[\alpha] \subset \mathbb{Z}[\alpha, t_1] \subset \mathbb{Z}[\alpha, t_1, t_2] \subset \dots$$

with the restriction maps $E_r : \mathbb{Z}[\alpha, t_1, \dots, t_{r+1}] \rightarrow \mathbb{Z}[\alpha, t_1, \dots, t_r]$

$$(E_r g)(\alpha, t_1, \dots, t_r) = g(\alpha, t_1, \dots, t_r, 0).$$

Since $f_{r+1}(\alpha, t_1, \dots, t_r, 0)(P^n) = f_r(\alpha, t_1, \dots, t_r)(P^n)$, we obtain the ring homomorphism

$$f : \mathcal{P} \rightarrow \text{Qsym}[t_1, t_2, \dots][\alpha] \subset \varprojlim_r \mathbb{Z}[\alpha, t_1, \dots, t_r] :$$

$$f(\alpha, t_1, t_2, \dots)(P^n) = \alpha^n + \sum_{k=1}^n \sum_{0 \leq a_1 < \dots < a_k \leq n-1} f_{a_1, \dots, a_k} \alpha^{a_1} M_{(n-a_k, \dots, a_2-a_1)}.$$

Definition 6.2. For a set $S = \{a_1, \dots, a_k\} \subset \{0, 1, \dots, n-1\}$ let us denote by $\omega(S)$ the composition $(n-a_k, a_k - a_{k-1}, \dots, a_2 - a_1)$ of the number $n - a_1$.

Then we can write

$$f(P^n) = \sum_{k \geq 0} \sum_{S: l(S)=k} f_S \alpha^{a_1} M_{\omega(S)}$$

6.2 Image

It follows from the formula, that the restriction

$$f(\alpha, t_1, \dots, t_r, t_{r+1}, \dots) \rightarrow f(\alpha, t_1, \dots, t_r, 0, 0, \dots) = f_r(\alpha, t_1, \dots, t_r)$$

is injective on the space of all n -dimensional polytopes, $n \leq r$.

Theorem 6.3. *The image of the space $\mathcal{P}^{2n}$ generated by all n -dimensional polytopes in the ring $\text{Qsym}[t_1, \dots, t_r][\alpha]$, $r \geq n$ under the mapping f_r consists of all the homogeneous polynomials g of degree $2n$ satisfying the equations*

1.

$$\begin{aligned} g(\alpha, t_1, -t_1, t_3, \dots, t_r) &= g(\alpha, 0, 0, t_3, \dots, t_r); \\ g(\alpha, t_1, t_2, -t_2, t_4, \dots, t_r) &= g(\alpha, t_1, 0, 0, t_4, \dots, t_r); \\ &\dots \\ g(\alpha, t_1, \dots, t_{r-2}, t_{r-1}, -t_{r-1}) &= g(\alpha, t_1, \dots, t_{r-2}, 0, 0); \end{aligned}$$

2.

$$g(-\alpha, t_1, \dots, t_{r-1}, \alpha) = g(\alpha, t_1, \dots, t_{r-1}, 0);$$

These equations are equivalent to the Bayer-Billera (generalized Dehn-Sommerville) relations.

Proof. $\Phi(-t)\Phi(t) = 1 = \Phi(0)\Phi(0)$, therefore

$$\begin{aligned} \xi_\alpha \Phi(t_r) \dots \Phi(t_3) \Phi(-t_1) \Phi(t_1) P^n &= \xi_\alpha \Phi(t_r) \dots \Phi(t_3) \Phi(0) \Phi(0) P^n; \\ \xi_\alpha \Phi(t_r) \dots \Phi(t_4) \Phi(-t_2) \Phi(t_2) \Phi(t_1) P^n &= \xi_\alpha \Phi(t_r) \dots \Phi(t_4) \Phi(0) \Phi(0) \Phi(t_1) P^n; \\ &\dots \\ \xi_\alpha \Phi(-t_{r-1}) \Phi(t_{r-1}) \Phi(t_{r-2}) \dots \Phi(t_1) P^n &= \xi_\alpha \Phi(0) \Phi(0) \Phi(t_{r-2}) \dots \Phi(t_1) P^n. \end{aligned}$$

On the other hand, Proposition 2.53 gives the last relation

$$\xi_{-\alpha} \Phi(\alpha) \Phi(t_{r-1}) \dots \Phi(t_1) P^n = \xi_\alpha \Phi(0) \Phi(t_{r-1}) \dots \Phi(t_1) P^n.$$

Now let us proof the opposite inclusion, that is if the homogeneous polynomial g of degree $2n$ satisfies the conditions of the theorem, then $g = f_r(p^n)$ for some $p^n \in \mathcal{P}^{2n}$.

Lemma 6.4. *Let P^n be an n -dimensional polytope. Then*

1. *The equation*

$$f_r(\alpha, t_1, \dots, t_q, -t_q, \dots, t_r) = f_r(\alpha, t_1, \dots, 0, 0, \dots, t_r)$$

is equivalent to the generalized Dehn-Sommerville relations

$$\sum_{j=a_t+1}^{a_{t+1}-1} (-1)^{j-a_t-1} f_{a_1, \dots, a_t, j, a_{t+1}, \dots, a_k} = (1 + (-1)^{a_{t+1}-a_t}) f_{a_1, \dots, a_t, a_{t+1}, \dots, a_k}$$

for $1 \leq k \leq \min\{r-1, n-1\}$, $k+1-q \leq t \leq r-q$.

2. *The equation*

$$f_r(-\alpha, t_1, \dots, t_{r-1}, \alpha) = f_r(\alpha, t_1, \dots, t_{r-1}, 0)$$

is equivalent to the generalized Dehn-Sommerville relations

$$\sum_{j=0}^{a_1-1} (-1)^j f_{j, a_1, \dots, a_k} = (1 + (-1)^{a_1-1}) f_{a_1, \dots, a_k}$$

for $0 \leq k \leq \min\{r-1, n-1\}$.

Proof.

$$\begin{aligned}
f_r(\alpha, t_1, \dots, t_q, -t_q, \dots, t_r) = \\
\alpha^n + \sum_{k=1}^{\min\{r, n\}} \sum_{0 \leq a_1 < \dots < a_k \leq n-1} f_{a_1, \dots, a_k} \alpha^{a_1} \left(\sum_{1 \leq l_1 < \dots < l_k \leq r} t_{l_1}^{n-a_k} \dots t_{l_k}^{a_2-a_1} \right) \Big|_{t_{q+1}=-t_q} = \\
= \alpha^n + \sum_{k=1}^{\min\{r-2, n\}} \sum_{0 \leq a_1 < \dots < a_k \leq n-1} f_{a_1, \dots, a_k} \alpha^{a_1} \left(\sum_{1 \leq l_1 < \dots < l_k \leq r, l_j \neq q, q+1} t_{l_1}^{n-a_k} \dots t_{l_k}^{a_2-a_1} \right) + \\
+ \sum_{k=1}^{\min\{r-1, n\}} \sum_{0 \leq a_1 < \dots < a_k \leq n-1} f_{a_1, \dots, a_k} \alpha^{a_1} \left(\sum_{1 \leq l_1 < \dots < l_j=q < \dots < l_k \leq r, l_{j+1} \neq q+1} t_{l_1}^{n-a_k} \dots t_q^{a_{k+2-j}-a_{k+1-j}} \dots t_{l_k}^{a_2-a_1} \right) + \\
+ \sum_{k=1}^{\min\{r-1, n\}} \sum_{0 \leq a_1 < \dots < a_k \leq n-1} f_{a_1, \dots, a_k} \alpha^{a_1} \left(\sum_{1 \leq l_1 < \dots < l_{j+1}=q+1 < \dots < l_k \leq r, l_j \neq q} t_{l_1}^{n-a_k} \dots (-t_q)^{a_{k+1-j}-a_{k-j}} \dots t_{l_k}^{a_2-a_1} \right) + \\
\sum_{k=1}^{\min\{r, n\}} \sum_{0 \leq a_1 < \dots < a_k \leq n-1} f_{a_1, \dots, a_k} \alpha^{a_1} \left(\sum_{1 \leq l_1 < \dots < l_j=q < l_{j+1}=q+1 < \dots < l_k \leq r} t_{l_1}^{n-a_k} \dots (-1)^{a_{k+1-j}-a_{k-j}} t_q^{a_{k+2-j}-a_{k-j}} \dots t_{l_k}^{a_2-a_1} \right)
\end{aligned}$$

The first and the second summands form exactly $f(\alpha, t_1, \dots, 0, 0, \dots, t_r)$. Therefore all the coefficients of the polynomial consisting of the last three summands should be equal to 0.

Consider the monomial $\alpha^{a_1} t_{l_1}^{n-a_k} \dots t_q^{a_{k+1}-a_k} \dots t_{l_k}^{a_2-a_1}$. Here $q = l_{k+1}-t$. The existence of a monomial of this form in the sum is equivalent to the conditions

$$k \leq \min\{r-1, n-1\}, \quad k+1-t \leq q, \quad t-1 \leq r-q-1.$$

The coefficient of the monomial should be equal to 0. This is equivalent to the relation:

$$(1 + (-1)^{a_{t+1}-a_t}) f_{a_1, \dots, a_t, a_{t+1}, \dots, a_k} + \sum_{j=a_t+1}^{a_{t+1}-1} (-1)^{j-a_t} f_{a_1, \dots, a_t, j, a_{t+1}, a_k} = 0$$

Now let us consider the remaining relation $f_r(-\alpha, t_1, \dots, t_{r-1}, \alpha) = f_r(\alpha, t_1, \dots, t_{r-1}, 0)$:

$$\begin{aligned}
&= (-\alpha)^n + \sum_{k=1}^{\min\{r-1, n\}} \sum_{0 \leq a_1 < \dots < a_k \leq n-1} f_{a_1, \dots, a_k} (-\alpha)^{a_1} \left(\sum_{1 \leq l_1 < \dots < l_k \leq r-1} t_{l_1}^{n-a_k} \dots t_{l_k}^{a_2-a_1} \right) + \\
&+ \sum_{k=1}^{\min\{r, n\}} \sum_{0 \leq a_1 < \dots < a_k \leq n-1} f_{a_1, \dots, a_k} (-\alpha)^{a_1} \left(\sum_{1 \leq l_1 < \dots < l_k=r} t_{l_1}^{n-a_k} \dots t_{l_{k-1}}^{a_3-a_2} \alpha^{a_2-a_1} \right) = \\
&\alpha^n + \sum_{k=1}^{\min\{r-1, n\}} \sum_{0 \leq a_1 < \dots < a_k \leq n-1} f_{a_1, \dots, a_k} \alpha^{a_1} \left(\sum_{1 \leq l_1 < \dots < l_k \leq r-1} t_{l_1}^{n-a_k} \dots t_{l_k}^{a_2-a_1} \right)
\end{aligned}$$

This is equivalent to the relations

$$(-1)^{a_1} f_{a_1, \dots, a_k} + \sum_{j=0}^{a_1-1} (-1)^j f_{j, a_1, \dots, a_k} = f_{a_1, \dots, a_k}$$

for $0 \leq k \leq \min\{r-1, n-1\}$. If $k=0$, then the corresponding relation is exactly the Euler formula

$$(-1)^n + (-1)^{n-1} f_{n-1} + \dots + f_2 - f_1 + f_0 = 1.$$

□

Corollary 6.5. For $r \geq n$ relations 1. and 2. of the theorem are equivalent to the generalized Dehn-Sommerville relations for the polytope P^n : For $S \subset \{0, \dots, n-1\}$, and $\{i, k\} \subseteq S \cup \{-1, n\}$ such that $i < k-1$ and $S \cap \{i+1, \dots, k-1\} = \emptyset$:

$$\sum_{j=i+1}^{k-1} (-1)^{j-i-1} f_{S \cup \{j\}} = (1 - (-1)^{k-i-1}) f_S.$$

Proof. We see that relations 1. and 2. follow from the generalized Dehn-Sommerville relations.

On the other hand, if $i = -1$ then the corresponding relation follows from equation 2.

If $S = \{a_1, \dots, a_t, a_{t+1}, \dots, a_s\}$, $i = a_t \geq 0$, and $k = a_{t+1}$; or $S = \{a_1, \dots, a_t\}$, $i = a_t$, and $k = n$, then we can take q such that $s+1-t \leq q \leq r-t$. $\square$

Remark 6.6. For n -dimensional polytopes for different $r \geq n$ not all the equations are independent.

For $i = -1$ relation (3) follows from equation 2. and all the equations of type 1. do not contain the case $i = -1$.

On the other hand, let $S = \{a_1, \dots, a_t, a_{t+1}, \dots, a_s\}$, $i = a_t \geq 0$, and $k = a_{t+1}$; or $S = \{a_1, \dots, a_t\}$, $i = a_t$, and $k = n$. Lemma 6.4 implies that relation (3) follows from the equation with $s+1-t \leq q \leq r-t$. Let us denote $a = s+1-t$, $b = r-t$. There are two conditions for s, t , namely

$$1 \leq s \leq n-1 \text{ and } 1 \leq t \leq s.$$

Let us rewrite these conditions in terms of a and b : $t = r-b$; $s = a+t-1 = a-b+r-1$, so

$$r-n \leq b-a \leq r-2; \quad b \leq r-1; \quad 1 \leq a$$

This gives us a triangle on the plane (a, b) . Each point (a, b) of this triangle corresponds to the conditions $a \leq q \leq b$, that is there should be the equation for q in the segment $[a, b]$. We can imagine that this segment is the segment $[(a, a), (a, b)]$ on the plane.

Thus for $r = n$ all the equations for $t_1, \dots, t_{n-1}$ are necessary, for $r = n+1$ it is enough to take the equations for $q = 2, 4, 6, \dots, 2[\frac{n}{2}]$. If $r \geq 2n-2$ one equation

$$f(\alpha, t_1, \dots, t_{n-1}, -t_{n-1}, t_{n+1}, \dots, t_r) = f(\alpha, t_1, \dots, 0, 0, t_{n+1}, \dots, t_r)$$

gives all the relations of type 1.

Now let us finish the proof of the theorem. If the homogeneous polynomial $g \in \text{Qsym}[t_1, \dots, t_r][\alpha]$ of degree $2n$ satisfies all the relations of the theorem, then its coefficients satisfy the generalized Dehn-Sommerville relations. Therefore all the coefficients are linear combinations of the coefficients $g_{a_1, \dots, a_k}$, where $S = \{a_1, \dots, a_k\} \in \Psi^n$. As we know, the vectors $\{f_S(Q), S \in \Psi^n\}$, $Q \in \Omega^n$ form a basis of the abelian group of all the vectors $\{f_S, S \in \Psi^n, f_S \in \mathbb{Z}\} = \mathbb{Z}^{c_n}$. So the vector $\{g_S, S \in \Psi^n\}$ is an integer combination of the vectors $\{f_S(Q), S \in \Psi^n\}$, $Q \in \Omega^n$. This implies that the polynomial g is an integer combination of the polynomials $f_r(Q)$, $Q \in \Omega^n$ with the same coefficients. $\square$

Let us remind that rank of the space $f_r(\mathcal{P}^{2n})$, $r \geq n$ is equal to c_n .

Proposition 6.7. Let $r \geq 2$, and let P^n be an n -dimensional polytope. Then

$$f_r(\alpha, t_1, \dots, t_r)(P^n) = f_1(\alpha, t_1 + \dots + t_r)(P^n)$$

if and only if P^n is simple. Here $f_1(\alpha, t)$ is a usual homogeneous f -polynomial in two variables.

Proof. On the ring of simple polytopes $d_k|_{\mathcal{P}_s} = \frac{d^k}{k!}|_{\mathcal{P}_s}$, so $\Phi(t) = e^{td}$.

Then $\Phi(t_r)\Phi(t_{r-1})\dots\Phi(t_1) = \Phi(t_1+\dots+t_r)$, therefore $f_r(\alpha, t_1, \dots, t_r)(P^n) = f_1(\alpha, t_1+\dots+t_r)(P^n)$.

On the other hand, let $f_r(\alpha, t_1, \dots, t_r)(P^n) = f_1(\alpha, t_1+\dots+t_r)(P^n)$.

Then $f_2(\alpha, t_1, t_2)(P^n) = f_1(\alpha, t_1+t_2)(P^n)$. So

$$\alpha^n + \sum_{i=0}^{n-1} f_i \alpha^i (t_1^{n-i} + t_2^{n-i}) + \sum_{0 \leq i < j \leq n-1} f_{ij} \alpha^i t_1^{n-j} t_2^{j-i} = \alpha^n + \sum_{i=0}^{n-1} f_i \alpha^i (t_1 + t_2)^{n-i}$$

In particular, $f_{01} = n f_0$, since the coefficients of the monomial $t_1^{n-1} t_2$ on the left and on the right are equal. Hence $2f_1 = f_{01} = n f_0$. This implies that the polytope P^n is simple. $\square$

Remark 6.8. Letting r tend to infinity we obtain that $f(\alpha, t_1, t_2, \dots)(P^n) = f_1(\alpha, t_1 + t_2 + \dots)(P^n)$ if and only if P^n is simple.

In the case of simple polytopes the equations of the first type are trivial, but the equation of the second type has the form:

$$f_1(-\alpha, t_1 + \dots + t_{r-1} + \alpha) = f_1(\alpha, t_1 + \dots + t_{r-1})$$

If we denote $t = t_1 + \dots + t_{r-1}$, then $f_1(-\alpha, \alpha + t) = f_1(\alpha, t)$. This equation is equivalent to the Dehn-Sommerville relations (after the change of variables $h(\alpha, t) = f_1(\alpha - t, t)$ it is equivalent to the fact that $h(\alpha, t) = h(t, \alpha)$)

6.3 Characterization of the generalized f -polynomial.

In this part we find the condition that uniquely determines the generalized f -polynomial.

At first let us find the relation between $f(P^n)$ and $f(d_k P^n)$.

Proposition 6.9. *For any polytope $P^n \in \mathcal{P}$ we have*

$$f(\alpha, t_1, t_2, \dots)(d_k P^n) = \frac{1}{k!} \frac{\partial^k}{\partial t^k} \Big|_{t=0} f(\alpha, t, t_1, t_2, \dots)(P^n)$$

Proof. Indeed, both expressions belong to $\mathcal{Qsym}[\alpha]$, and they are uniquely defined by their restrictions to n variables. Then we have

$$\begin{aligned} \left(\frac{1}{k!} \frac{\partial^k}{\partial t^k} \Big|_{t=0} f_{n+1}(\alpha, t, t_1, t_2, \dots)(P^n) \right) \Big|_{t_{n+1}=t_{n+2}=\dots=0} &= \\ \frac{1}{k!} \frac{\partial^k}{\partial t^k} \Big|_{t=0} f_{n+1}(\alpha, t, t_1, t_2, \dots, t_n)(P^n) &= \frac{1}{k!} \frac{\partial^k}{\partial t^k} \Big|_{t=0} \xi_\alpha \Phi(t_n) \dots \Phi(t_1) \Phi(t) P^n = \xi_\alpha \Phi(t_n) \dots \Phi(t_1) d_k P^n = \\ f_n(\alpha, t_1, t_2, \dots, t_n)(d_k P^n) &= f(\alpha, t_1, t_2, \dots)(d_k P^n) \Big|_{t_{n+1}=t_{n+2}=\dots=0} \end{aligned}$$

□

Corollary 6.10.

$$\begin{aligned} f(\alpha, t, t_1, t_2, \dots)(P^n) &= f(\alpha, t_1, t_2, \dots)(P^n) + f(\alpha, t_1, t_2, \dots)(dP^n)t + \\ &\quad + f(\alpha, t_1, t_2, \dots)(d_2 P^n)t^2 + \dots + f(\alpha, t_1, t_2, \dots)(d_n P^n)t^n \\ f_{r+1}(\alpha, t, t_1, \dots, t_r)(P^n) &= f_r(\alpha, t_1, \dots, t_r)(P^n) + f_r(\alpha, t_1, \dots, t_r)(dP^n)t + \\ &\quad + f_r(\alpha, t_1, \dots, t_r)(d_2 P^n)t^2 + \dots + f_r(\alpha, t_1, \dots, t_r)(d_n P^n)t^n, \quad r \geq 0; \end{aligned}$$

Proof. The first equality follows from Proposition 6.9.

In fact, both equalities can be proved directly:

$$\begin{aligned} f_{r+1}(\alpha, t, t_1, \dots, t_r)(P^n) &= \xi_\alpha \Phi(t_r) \dots \Phi(t_1) \Phi(t) P^n = \sum_{k=0}^n (\xi_\alpha \Phi(t_r) \dots \Phi(t_1) d_k P^n) t^k = \\ &= \sum_{k=0}^n f_r(\alpha, t_1, \dots, t_r)(d_k P^n) t^k. \end{aligned}$$

Letting r tend to infinity we obtain the first equality. □

The first equality of Corollary 6.10 is equivalent to the condition $f(\alpha, t_1, t_2, \dots)(\Phi(t)P^n) = f(\alpha, t, t_1, t_2, \dots)(P^n)$.

Thus we see that the following diagram commutes

$$\begin{array}{ccc} \mathcal{P} & \xrightarrow{f} & \text{Qsym}[t_1, t_2, \dots][\alpha] \\ \Phi \downarrow & & T \downarrow \\ \mathcal{P}[t] & \xrightarrow{f} & \text{Qsym}[t, t_1, t_2, \dots][\alpha] \end{array}$$

where $f(t) = t$, and $T : \text{Qsym}[t_1, t_2, \dots][\alpha] \rightarrow \text{Qsym}[t, t_1, t_2, \dots][\alpha]$ is a ring homomorphism:

$$Tg(\alpha, t_1, t_2, \dots) = g(\alpha, t, t_1, t_2, \dots), \quad g \in \text{Qsym}[t_1, t_2, \dots][\alpha].$$

Remark 6.11. Note that $f_r(\alpha, t_1, \dots, t_r)(\Phi(t)P^n) = f_{r+1}(\alpha, t, t_1, \dots, t_r)(P^n)$ for all $r \geq 0$.

Consider the ring homomorphism $T_{r+1} : \text{Qsym}[t_1, \dots, t_r][\alpha] \rightarrow \text{Qsym}[t, t_1, \dots, t_r][\alpha]$ defined as

$$T_{r+1}(\alpha) = \alpha, \quad T_{r+1}M_\omega(t_1, \dots, t_r) = M_\omega(t, t_1, \dots, t_r)$$

Then the corresponding diagram $f_r(\alpha, t_1, \dots, t_r)(\Phi(t)P^n) = (T_{r+1}f_r)(\alpha, t, t_1, \dots, t_r)$ commutes only for $r \geq n$.

Theorem 6.12. Let $\psi : \mathcal{P} \rightarrow \text{Qsym}[t_1, t_2, \dots][\alpha]$ be a linear map such that

1. $\psi(\alpha, 0, 0, \dots)(P^n) = \alpha^n$;
2. The following diagram commutes:

$$\begin{array}{ccc} \mathcal{P} & \xrightarrow{\psi} & \text{Qsym}[t_1, t_2, \dots][\alpha] \\ \Phi \downarrow & & T \downarrow \\ \mathcal{P}[t] & \xrightarrow{\psi} & \text{Qsym}[t, t_1, t_2, \dots][\alpha] \end{array}$$

Then $\psi = f$.

Proof. The first condition implies that $\psi(\alpha, 0, 0, \dots)(p) = \xi_\alpha p$ for any $p \in \mathcal{P}$.

Let $\psi(P^n) = \alpha^n + \sum_{\omega} \psi_\omega(\alpha)M_\omega$, and let $\omega = (j_1, \dots, j_k)$.

Since $\psi(\alpha, t, t_1, t_2, \dots)(P^n) = \psi(\alpha, t_1, t_2, \dots)(\Phi(t)P^n)$, we have

$$\psi(\alpha, y_1, y_2, \dots, y_k, t_1, t_2, \dots)(P^n) = \psi(\alpha, t_1, t_2, \dots)(\Phi(y_k) \dots \Phi(y_1)P^n)$$

Therefore,

$$\psi(\alpha, y_1, \dots, y_k, 0, 0, \dots) = \psi(\alpha, 0, 0, \dots)(\Phi(y_k) \dots \Phi(y_1)P^n) = \xi_\alpha \Phi(y_k) \dots \Phi(y_1)P^n = f(\alpha, y_1, \dots, y_k, 0, 0, \dots)$$

Hence $\psi_\omega(\alpha) = f_{S(\omega)}\alpha^{n-|\omega|}$, where $S(\omega) = \{n - |\omega|, n - |\omega| + j_k, \dots, n - j_1\}$. This is valid for all ω , so $\psi = f$. $\square$

Remark 6.13. Let us mention that the mapping T is an isomorphism $\text{Qsym}[t_1, t_2, \dots][\alpha] \rightarrow \text{Qsym}[t, t_1, t_2, \dots][\alpha]$, while Φ is an injection and its image is described by the condition: $p(t) \in \Phi(\mathcal{P})$ is and only if $\Phi(-t)p(t) \in \mathcal{P}$.

On the ring of simple polytopes we have $f(\alpha, t_1, t_2, \dots)(p) = f_1(\alpha, t_1 + t_2 + \dots)(p) = f_1(\alpha, \sigma_1)(p)$, so the image of $\mathcal{P}_s$ belongs to $\mathbb{Z}[\alpha, \sigma_1]$. On the other hand, $\Phi(t) = e^{td}$, so we have the condition $f_1(\alpha, t_1)(e^{td}p) = f_1(\alpha, t + t_1)(p)$.

Proposition 6.14. Let $\psi : \mathcal{P}_s \rightarrow \mathbb{Z}[\alpha, t]$ be a linear mapping such that

1. $\psi(\alpha, 0)(P^n) = \alpha^n$;
2. One of the following equivalent conditions holds:

$$(a) \quad \psi(\alpha, t_1)(e^{td}p) = \psi(\alpha, t + t_1)(p);$$

$$(b) \quad \psi(\alpha, t)(dp) = \frac{\partial}{\partial t} \psi(\alpha, t)(p).$$

Then $\psi = f_1$.

Proof. Let conditions 1 and 2a hold. Then

$$\psi(\alpha, t + t_1)(p) = \psi(\alpha, t_1)(e^{td}p) = \psi(\alpha, 0)(e^{(t+t_1)d}p) = \xi_\alpha \Phi(t + t_1)p = f_1(\alpha, t_1 + t)(p).$$

It remains to prove that conditions 2a and 2b are equivalent.

Indeed, if $\psi(\alpha, t_1)(e^{td}p) = \psi(\alpha, t + t_1)(p)$, then $\psi(\alpha, t)(p) = \psi(\alpha, 0)(e^{td}p)$, therefore

$$\frac{\partial}{\partial t} \psi(\alpha, t)(p) = \frac{\partial}{\partial t} \psi(\alpha, 0)(e^{td}p) = \psi(\alpha, 0)(e^{td}dp) = \psi(\alpha, t)(dp).$$

Now let $\psi(\alpha, t)(dp) = \frac{\partial}{\partial t} \psi(\alpha, t)(p)$. Then

$$\frac{\partial}{\partial t} \psi(\alpha, t_1)(e^{td}p) = \psi(\alpha, t_1)(d(e^{td}p)) = \frac{\partial}{\partial t_1} \psi(\alpha, t_1)(e^{td}p)$$

Hence $\left(\frac{\partial}{\partial t} - \frac{\partial}{\partial t_1}\right) \psi(\alpha, t_1)(e^{td}p) = 0$, so $\psi(\alpha, t_1)(e^{td}p)$ depends on α and $t + t_1$.

Thus $\psi(\alpha, t_1)(e^{td}p) = \psi(\alpha, t + t_1)(e^{0d}p) = \psi(\alpha, t + t_1)(p)$. $\square$

Proposition 6.14 in the form 1, 2b was first proved in [Buch].

7 Structure of $\mathcal{D}^*$

7.1 Main theorem

Definition 7.1. Let us denote by $\mathcal{D}^*$ the graded dual Hopf algebra to $\mathcal{D}$.

Let us identify the Hopf algebra $\mathcal{M}$, which is the graded dual $\mathcal{Z}^*$ to $\mathcal{Z}$, with $\text{Qsym}[t_1, t_2, \dots]$.

The mappings $\mathcal{L}: \mathcal{Z} \rightarrow \mathcal{D}$, $\mathcal{R}: \mathcal{Z}^{op} \rightarrow \mathcal{D}$, $Z_i \rightarrow d_i$ are surjections, so the dual mappings are injections. So the ring homomorphisms $\mathcal{L}^*$, and $\mathcal{R}^* = \varrho^* \circ \mathcal{L}^*$ define two embeddings $\mathcal{D}^* \subset \text{Qsym}$, which are related by the involution $\varrho^* = *$.

In fact, $\mathcal{L}^*$ defines an embedding $\mathcal{D}^* \subset \text{Qsym}$ as a Hopf subalgebra, while for $\mathcal{R}^*$ we have

$$\Delta \circ \mathcal{R}^* = \tau_{\text{Qsym}} \circ (\mathcal{R}^* \otimes \mathcal{R}^*) \circ \Delta,$$

where τ_{Qsym} is a homomorphism $\text{Qsym} \otimes \text{Qsym} \rightarrow \text{Qsym} \otimes \text{Qsym}$ that interchanges the tensor factors: $\tau_{\text{Qsym}}(x \otimes y) = y \otimes x$.

Definition 7.2. For each r let us define the ring homomorphism $\mathcal{R}_r^*: \mathcal{D}^* \rightarrow \text{Qsym}[t_1, \dots, t_r]$ by the formula

$$\mathcal{R}_r^*(\psi) = \mathcal{R}^*(\psi)(t_1, \dots, t_r, 0, 0, \dots)$$

Since the restriction $\text{Qsym} \rightarrow \text{Qsym}[t_1, \dots, t_r]$ is injective for any graduation $(2n)$, $n \leq r$, we see that $\mathcal{R}_r^*$ is injective for graduations $(2n)$, $n \leq r$, of $\mathcal{D}^*$.

Theorem 7.3. Let $r \geq n$. Then the image of the $(2n)$ -th, $n \geq 1$, graded component of the ring $\mathcal{D}^*$ in the ring $\text{Qsym}[t_1, \dots, t_r]$ under the map $\mathcal{R}_r^*$ consists of all the homogeneous polynomials of degree $2n$ satisfying the relations:

$$\begin{aligned} g(t_1, -t_1, t_3, \dots, t_r) &= g(0, 0, t_3, \dots, t_r); \\ g(t_1, t_2, -t_2, t_4, \dots, t_r) &= g(t_1, 0, 0, t_4, \dots, t_r); \\ &\dots \\ g(t_1, \dots, t_{r-2}, t_{r-1}, -t_{r-1}) &= g(t_1, \dots, t_{r-2}, 0, 0). \end{aligned}$$

Proof. The ring $\mathcal{R}^*(\mathcal{D}^*) \subset \mathcal{Z}^*$ consists of all the linear functions $\psi \in \mathcal{Z}^*$ satisfying the property:

$$\langle \psi, z_1 \Phi(t) \Phi(-t) z_2 \rangle = \langle \psi, z_1 z_2 \rangle \quad (15)$$

for all $z_1, z_2 \in \mathcal{Z}$, that is the coefficients of all t^k , $k \geq 1$, on the left are equal to 0.

We have

$$\mathcal{R}^*(\psi) = \sum_{\omega} \langle \mathcal{R}^*(\psi), Z_{\omega} \rangle = \sum_{\omega} \langle \psi, \mathcal{R} Z_{\omega} \rangle M_{\omega}$$

Therefore,

$$\begin{aligned} \mathcal{R}_r^*(\psi) &= \sum_{k=0}^r \sum_{(j_1, \dots, j_k)} \langle \psi, d_{j_k} \dots d_{j_1} \rangle M_{(j_1, \dots, j_k)}(t_1, \dots, t_r) = \\ &= \sum_{k=0}^r \sum_{(j_1, \dots, j_k)} \langle \psi, d_{j_k} \dots d_{j_1} \rangle \sum_{1 \leq l_1 < \dots < l_k \leq r} t_{l_1}^{j_1} \dots t_{l_k}^{j_k} = \\ &= \sum_{k=0}^r \sum_{(j_1, \dots, j_k)} \sum_{1 \leq l_1 < \dots < l_k \leq r} \langle \psi, d_{j_k} t_{l_k}^{j_k} \dots d_{j_1} t_{l_1}^{j_1} \rangle = \langle \psi, \Phi(t_r) \dots \Phi(t_1) \rangle \end{aligned}$$

Then

$$\begin{aligned} \mathcal{R}_r^*(\psi)(t_1, \dots, t_i, -t_i, \dots, t_r) &= \langle \psi, \Phi(t_r) \dots \Phi(-t_i) \Phi(t_i) \dots \Phi(t_1) \rangle = \\ &= \langle \psi, \Phi(t_r) \dots \Phi(0) \Phi(0) \dots \Phi(t_1) \rangle = \mathcal{R}_r^*(\psi)(t_1, \dots, 0, 0, \dots, t_r) \end{aligned}$$

On the other hand, let $g \in \text{Qsym}[t_1, \dots, t_r]$ be a homogeneous polynomial of degree $2n$ satisfying all the relations of the theorem. Consider a unique homogeneous function $\psi \in \mathcal{M}$ such that $\deg \psi = 2n$, and $\psi(t_1, \dots, t_r, 0, 0, \dots) = g$. Let us prove that ψ belongs to the image of $\mathcal{D}^*$ under the embedding $\mathcal{R}^*$.

It is sufficient to prove the relation (15) in the case when z_1, z_2 are monomials. Since

$$\Phi(t) \Phi(-t) = 1 + (Z_1 - Z_1)t + (2Z_2 - Z_1^2)t^2 + \dots = 1 + (2Z_2 - Z_1^2)t^2 + \dots, \text{ and } \deg \psi = 2n,$$

the cases $\deg z_1 + \deg z_2 = 2n$ and $2(n-1)$ are trivial. Let

$$z_1 = Z_{\omega}, \quad z_2 = Z_{\omega'}, \quad \omega = (j_1, \dots, j_l), \quad \omega' = (j'_1, \dots, j'_{l'}), \quad |\omega| + |\omega'| = n - k, \quad k \geq 2.$$

Then the only equality we need to prove is

$$\langle \psi, Z_{\omega} \left(\sum_{i=0}^k (-1)^i Z_{k-i} Z_i \right) Z_{\omega'} \rangle = 0$$

Let us consider the equality

$$g(t_1, \dots, t_l, t_{l+1}, -t_{l+1}, t_{l+3}, \dots, t_{l+2+l'}, \dots, t_r) = g(t_1, \dots, t_l, 0, 0, t_{l+3}, \dots, t_{l+2+l'}, \dots, t_r).$$

The coefficient of the monomial $t_1^{j_1} \dots t_l^{j_l} t_{l+1}^k t_{l+3}^{j'_1} \dots t_{l+2+l'}^{j'_{l'}}$ on the left is exactly

$$\sum_{i=0}^k (-1)^i \langle \psi, Z_{\omega} Z_{k-i} Z_i Z_{\omega'} \rangle = \langle \psi, Z_{\omega} \left(\sum_{i=0}^k (-1)^i Z_{k-i} Z_i \right) Z_{\omega'} \rangle,$$

and on the right it is equal to 0. So $\langle \psi, Z_{\omega} \Phi(t) \Phi(-t) Z_{\omega'} \rangle = 0$ for all ω, ω' , therefore, $\psi \in \mathcal{R}^*(\mathcal{D}^*)$. $\square$

Definition 7.4. Let us define the operations $\Theta_k : \mathbb{Z}[t_1, t_2, \dots] \rightarrow \mathbb{Z}[t_1, t_2, \dots]$ as

$$\Theta_k g(t_1, t_2, \dots) = g(t_1, \dots, t_{k-1}, t, -t, t_k, t_{k+1}, \dots)$$

Corollary 7.5. The ring $\mathcal{R}^*(\mathcal{D}^*) \subset \mathcal{Qsym}$ is defined by the equations

$$g \in \mathcal{R}^*(\mathcal{D}^*) \text{ if and only if } \Theta_k g = g \text{ for all } k \geq 1.$$

Proof. Let g be a homogenous quasi-symmetric function of degree $2n$ representing the function $\psi \in \mathcal{D}^*$. Since g has bounded degree, each monomial of $g(t_1, \dots, t_{k-1}, t, -t, t_k, t_{k+1}, \dots)$ appears with the same coefficient in the polynomial $g_{r+2}(t_1, \dots, t_{k-1}, t, -t, t_k, \dots, t_r) = g(t_1, \dots, t_{k-1}, t, -t, t_k, \dots, t_r, 0, 0, \dots)$, when r is large enough. Then Theorem 7.3 implies that

$$g_{r+2}(t_1, \dots, t_{k-1}, t, -t, t_k, \dots, t_r) = g_{r+2}(t_1, \dots, t_{k-1}, 0, 0, t_k, \dots, t_r).$$

Therefore each monomial containing t has coefficient 0, so $g(t_1, \dots, t_{k-1}, t, -t, t_k, \dots)$ does not depend on t . So we can set $t = 0$ to obtain

$$g(t_1, \dots, t_{k-1}, t, -t, t_k, \dots) = g(t_1, \dots, t_{k-1}, 0, 0, t_k, \dots) = g(t_1, \dots, t_{k-1}, t_k, \dots)$$

On the other hand, let $g \in \mathcal{Qsym}^{2n}$ be a quasi-symmetric function such that $\Theta_k g = g$ for all $k \geq 1$. Then for $g_n(t_1, \dots, t_n) = g(t_1, \dots, t_n, 0, 0, \dots)$ we have

$$\begin{aligned} g_n(t_1, \dots, t_{i-1}, t_i, -t_i, t_{i+2}, \dots, t_n) &= g(t_1, \dots, t_{i-1}, t_i, -t_i, t_{i+2}, \dots, t_n, 0, 0, \dots) = \\ &= g(t_1, \dots, t_{i-1}, t_{i+2}, \dots, t_n, 0, 0, \dots) = g_n(t_1, \dots, t_{i-1}, t_{i+2}, \dots, t_n, 0, 0) = g_n(t_1, \dots, t_{i-1}, 0, 0, t_{i+2}, \dots, t_n) \end{aligned}$$

for $1 \leq i \leq n-1$. Then Theorem 7.3 implies that $g_n = \mathcal{R}_n^*(\psi)$ for some function $\psi \in \mathcal{D}^*$ of degree $2n$. Since the restriction $g(t_1, t_2, \dots) \rightarrow g(t_1, \dots, t_n, 0, 0, \dots)$ is injective for graduation $2n$, we have $g = \mathcal{R}^*\psi$. $\square$

Remark 7.6. Since $\mathcal{R}^* = \varrho^* \circ \mathcal{L}^*$, and the subring $\mathcal{R}^*(\mathcal{D}^*) \subset \mathcal{Qsym}$ is invariant under the involution $\varrho^* = *$, Theorem 7.3 and Corollary 7.5 remain valid for the mapping $\mathcal{L}^*$.

Proposition 7.7. $\mathcal{D}^* \otimes \mathbb{Q}$ is a free polynomial algebra

$$\mathcal{D}^* \otimes \mathbb{Q} \simeq \mathbb{Q}[\text{LYN}_{\text{odd}}],$$

where LYN_{odd} are Lyndon words consisting of odd positive integers. Rank of the $(2n)$ -th graded component of $\mathcal{D}^* \otimes \mathbb{Q}$ is equal to the $(n-1)$ -th Fibonacci number c_{n-1} .

Proof. This follows from Proposition 5.10, Corollary 3.21 to the shuffle algebra structure theorem, and Corollary 5.7. $\square$

7.2 Applications

Let $\mathcal{D}^*[\alpha]$ be a graded ring of polynomials in variable α , where $\deg \alpha = 2$, and for the element ψ_i of graduation $2i$ the element $\psi_i \alpha^j$ has graduation $2(i+j)$.

Let us consider the mapping $\varphi_\alpha: \mathcal{P} \rightarrow \mathcal{D}^*[\alpha]$, defined by the formula

$$\langle \varphi_\alpha(P), D \rangle = \xi_\alpha(DP)$$

for $D \in \mathcal{D}$. Then

$$\langle \varphi_\alpha(P \times Q), D \rangle = \xi_\alpha(D(P \times Q)) = \xi_\alpha \left(\sum_i D'_i P \times D''_i Q \right) = \sum_i (\xi_\alpha D'_i P) (\xi_\alpha D''_i Q) = \langle \varphi_\alpha(P) \cdot \varphi_\alpha(Q), D \rangle,$$

where $\Delta(D) = \sum_i D'_i \otimes D''_i$. So φ_α is a ring homomorphism.

Proposition 7.8. We have $\mathcal{R}^* \circ \varphi_\alpha = f$, where f is the generalized f -polynomial.

Proof. Indeed,

$$\begin{aligned} \mathcal{R}^* \circ \varphi_\alpha(P) &= \sum_\omega \langle \mathcal{R}^* \circ \varphi_\alpha(P), Z_\omega \rangle M_\omega = \sum_\omega \langle \varphi_\alpha(P), \mathcal{R} Z_\omega \rangle M_\omega = \\ &= \sum_\omega \langle \varphi_\alpha(P), D_{\omega^*} \rangle M_\omega = \sum_\omega \xi_\alpha(D_\omega P) M_{\omega^*} = f(P) \end{aligned}$$

$\square$

Proposition 7.9. *The image of the space $\mathcal{P}^{2n}$ under the mapping $\varphi_\alpha : \mathcal{P} \rightarrow \mathcal{D}^*[\alpha]$ consists of all homogeneous functions $\psi(\alpha)$ of degree $2n$ such that*

$$\langle \psi(-\alpha), \Phi(\alpha)D \rangle = \langle \psi(\alpha), D \rangle \quad (16)$$

for all $D \in \mathcal{D}$.

Proof. According to Proposition 2.53 we have $\xi_{-\alpha}\Phi(\alpha) = \xi_\alpha$, so if $\psi(\alpha) = \varphi_\alpha(p)$, then

$$\langle \psi(-\alpha), \Phi(\alpha)D \rangle = \xi_{-\alpha}\Phi(\alpha)Dp = \xi_\alpha Dp = \langle \psi(\alpha), D \rangle.$$

On the other hand, let condition (16) hold. Then the polynomial $g(\alpha, t_1, \dots, t_n) = \mathcal{R}_n^* \psi(\alpha)$ satisfies the relation

$$\begin{aligned} g(-\alpha, t_1, \dots, t_{n-1}, \alpha) &= \langle \psi(-\alpha), \Phi(\alpha)\Phi(t_{n-1}) \dots \Phi(t_1) \rangle = \\ &= \langle \psi(\alpha), \Phi(0)\Phi(t_{n-1}) \dots \Phi(t_1) \rangle = g(\alpha, t_1, \dots, t_{n-1}, 0). \end{aligned}$$

Theorem 7.3 implies that the polynomial g satisfies all the relations of Theorem 6.3, so $g = f_n(p^n)$ for some $p^n \in \mathcal{P}^{2n}$. We have $\mathcal{R}_n^* \psi(\alpha) = g = f_n(p^n) = \mathcal{R}_n^* \varphi_\alpha(p^n)$. Since the restriction $\mathcal{R}^* \rightarrow \mathcal{R}_n^*$ is injective on the $(2j)$ -th graded component of the ring $\mathcal{D}^*$ for $j \leq n$, and $\mathcal{R}^*$ is an embedding, we obtain that $\psi(\alpha) = \varphi_\alpha(p^n)$. $\square$

There is a right action of the ring $\mathcal{D}$ on its graded dual $\mathcal{D}^*$:

$$(\psi, D) \rightarrow \psi D: \quad \langle \psi D, D' \rangle = \langle \psi, DD' \rangle.$$

If $\deg \psi = 2n$ and $\deg D = 2k$, then $\deg \psi D = 2(n-k)$. Similarly, there is an action of $\mathcal{D}[[\alpha]]$ on $\mathcal{D}^*[[\alpha]]$. Then condition (16) means that $\psi(-\alpha)\Phi(\alpha) = \psi(\alpha)$. Since for any homogeneous function $\psi(\alpha)$ of degree $2n$ the function $\psi(\alpha)\Phi(\alpha)$ still has degree $2n$, we obtain the corollary.

Corollary 7.10. *Let $\psi(\alpha) \in \mathcal{D}^*[\alpha]$. Then*

$$\psi(\alpha) \in \varphi_\alpha(\mathcal{P}) \Leftrightarrow \psi(-\alpha)\Phi(\alpha) = \psi(\alpha). \quad (17)$$

Let $\psi(\alpha) = \psi_0 + \psi_1\alpha + \dots + \psi_n\alpha^n$, $u(\alpha) = \psi_0 + \psi_2\alpha^2 + \psi_4\alpha^4 + \dots$, $w(\alpha) = \psi_1\alpha + \psi_3\alpha^3 + \dots$. Then $u(\alpha)\Phi(\alpha) - w(\alpha)\Phi(\alpha) = u(\alpha) + w(\alpha)$. Therefore,

$$u(\alpha)(\Phi(\alpha) - 1) = w(\alpha)(\Phi(\alpha) + 1). \quad (18)$$

For example,

$$\psi_0 d = 2\psi_1; \quad \psi_0 d_3 + \psi_2 d = \psi_1 d_2 + 2\psi_3.$$

Consider the Hopf algebra $\mathcal{D} \otimes \mathbb{Z}[\frac{1}{2}]$. Then for the graded dual over $\mathbb{Z}[\frac{1}{2}]$ Hopf algebra we have: $(\mathcal{D} \otimes \mathbb{Z}[\frac{1}{2}])^* \simeq \mathcal{D}^* \otimes \mathbb{Z}[\frac{1}{2}]$. The condition that describes the image of the ring $\mathcal{P} \otimes \mathbb{Z}[\frac{1}{2}]$ in $\mathcal{D}^* \otimes \mathbb{Z}[\frac{1}{2}][\alpha]$ is the same, namely relation (18).

Proposition 7.11. *In the ring $\mathcal{D}^* \otimes \mathbb{Z}[\frac{1}{2}][\alpha]$ relation (18) is equivalent to the relation:*

$$w(\alpha) = u(\alpha) \frac{\Phi(\alpha) - 1}{\Phi(\alpha) + 1} = u(\alpha) \frac{\frac{\Phi(\alpha) - 1}{2}}{1 + \frac{\Phi(\alpha) - 1}{2}} = u(\alpha) \sum_{k=1}^{\infty} (-1)^{k-1} \left(\frac{\Phi(\alpha) - 1}{2} \right)^k. \quad (19)$$

Remark 7.12. In fact, equation (19) implies that the function $w(\alpha)$ is odd, if the function $u(\alpha)$ is even. Indeed,

$$\begin{aligned} w(-\alpha) &= u(-\alpha) \frac{\Phi(-\alpha) - 1}{\Phi(-\alpha) + 1} = u(\alpha) \frac{\frac{1}{\Phi(\alpha)} - 1}{\frac{1}{\Phi(\alpha)} + 1} = u(\alpha)(1 - \Phi(\alpha))\Phi(\alpha)^{-1}(1 + \Phi(\alpha))^{-1}\Phi(\alpha) = \\ &= -u(\alpha) \frac{\Phi(\alpha) - 1}{\Phi(\alpha) + 1} = -w(\alpha). \end{aligned}$$

Therefore for any even function $u(\alpha)$ the function $\psi(\alpha) = u(\alpha) + u(\alpha) \frac{\Phi(\alpha) - 1}{\Phi(\alpha) + 1} = u(\alpha) \frac{2\Phi(\alpha)}{\Phi(\alpha) + 1}$ belongs to the image of φ_α

Remark 7.13. The same relation is valid in $\mathcal{D}^* \otimes \mathbb{Q}[\alpha]$.

Let us take $\varphi_\alpha(p)$ with $\alpha = 0$. Then we obtain the classical map

$$\varphi_0 : \mathcal{P} \rightarrow \mathcal{D}^* : p \rightarrow \varphi_0(p) \quad \langle \varphi_0(p), D \rangle = \xi_0 D p, \quad \forall p \in \mathcal{P}, D \in \mathcal{D}$$

Proposition 7.14. *The mapping $\varphi_0 \otimes 1 : \mathcal{P} \otimes \mathbb{Z}[\frac{1}{2}] \rightarrow \mathcal{D}^* \otimes \mathbb{Z}[\frac{1}{2}]$ is a surjection.*

Proof. Let $\psi_0 \in \mathcal{D}^* \otimes \mathbb{Z}[\frac{1}{2}]$. Consider the element $\psi(\alpha) = u(\alpha) + w(\alpha)$, $u(\alpha) = \psi_0$, $w(\alpha) = \psi_0 \frac{\Phi(\alpha)-1}{\Phi(\alpha)+1}$. According to $\mathbb{Z}[\frac{1}{2}]$ -version of Corollary 7.10 we obtain that $\psi(\alpha) = \varphi_\alpha(p) \in \varphi_\alpha \otimes 1(\mathcal{P} \otimes \mathbb{Z}[\frac{1}{2}])$. Then $\psi_0 = \psi(0) = \varphi_0(p)$. $\square$

Question 7.15. What is the image of the map φ_0 over the integers?

Remark 7.16. Let us note, that the space $\mathcal{P}/\text{Ker } \varphi_\alpha$ consists of the equivalence classes of integer combinations of polytopes under the equivalence relation: $p \sim q$ if and only if p and q have equal flag f -vectors. Rank of the $(2n)$ -th graded component of the group $\mathcal{P}/\text{Ker } \varphi_\alpha$ is equal to c_n ([BB], see also Section 2.5).

On the other hand, by Corollary 5.7 rank of the $(2n)$ -th graded component of the ring $\mathcal{D}^*$ is equal to c_{n-1} , so the mapping $\varphi_0 : \mathcal{P}/\text{Ker } \varphi_\alpha \rightarrow \mathcal{D}^*$ is not injective.

Example 7.17. Let us consider small dimensions.

- $n = 1$. $\mathcal{P}/\text{Ker } \varphi_\alpha$ is generated by $CC = I$. The ring $\mathcal{D}$ in this graduation is generated by d . Then $\varphi_0(I) = 2d^*$.
- $n = 2$. $\mathcal{P}/\text{Ker } \varphi_\alpha$ is generated by $CCC = \Delta^2$ and $BCC = I^2$. $\mathcal{D}$ has one generator d_2 . Then $\varphi_0(\Delta^2) = 3d_2^*$, $\varphi_0(I^2) = 4d_2^*$. Therefore, $d_2^* = \varphi_0(I^2 - \Delta^2)$, $\text{Ker } \varphi_0$ is generated by $3I^2 - 4\Delta^2$.
- $n = 3$. $\mathcal{P}/\text{Ker } \varphi_\alpha$ is generated by $BCCC, CBCC = CI^2, CCCC = \Delta^3$, while $\mathcal{D}$ is generated by d_3, d_2d . Then

$$\varphi_0(BC^3) = 5d_3^* + 18(d_2d)^*, \quad \varphi_0(CBC^2) = 5d_3^* + 16(d_2d)^*, \quad \varphi_0(C^4) = 4d_3^* + 12(d_2d)^*$$

So $\text{Im } \varphi_0$ has the basis $d_3^*, 2(d_2d)^*$ and $\text{Ker } \varphi_0$ is generated by $2BC^3 - 6CBC^2 + 5C^4$.

Here by D_ω^* we denote the element of the dual basis: $\langle D_\omega^*, D_\sigma \rangle = \delta_{\omega, \sigma}$

8 Multiplicative Structure of $f(\mathcal{P}) \otimes \mathbb{Q}$

Theorem 8.1. *The ring $f(\mathcal{P}) \otimes \mathbb{Q}$ is a free polynomial algebra.*

Proof. According to Proposition 7.7 we have:

$$\mathcal{D}^* \otimes \mathbb{Q} \simeq \mathbb{Q}[\text{LYN}_{\text{odd}}],$$

where LYN_{odd} are Lyndon words consisting of odd positive integers. This is a free polynomial algebra. Therefore $\mathcal{D}^* \otimes \mathbb{Q}[\alpha]$ is a free polynomial algebra in the generators LYN_{odd} and α .

Let $\{f_\lambda\}$ be the functions in $\mathcal{D}^* \otimes \mathbb{Q}$ corresponding to the Lyndon words. Consider the polynomials $f_\lambda(\alpha)$ defined as

$$f_\lambda(\alpha) = u_\lambda(\alpha) + w_\lambda(\alpha); \quad u_\lambda(\alpha) = f_\lambda, \quad w_\lambda(\alpha) = f_\lambda \frac{\Phi(\alpha) - 1}{\Phi(\alpha) + 1} = f_\lambda \frac{e^{s(\alpha)} - 1}{e^{s(\alpha)} + 1}$$

Each polynomial $f_\lambda(\alpha)$ is a homogeneous element of degree $\deg f_\lambda$. Let us identify $\mathcal{D}^*$ with its image in $\text{Qsym}[t_1, t_2, \dots]$ under the embedding $\mathcal{R}^*$.

Then according to Remark 7.13 we have: $f_\lambda(\alpha) \in f(\mathcal{P} \otimes \mathbb{Q}) = f(\mathcal{P}) \otimes \mathbb{Q}$.

Lemma 8.2. $f(\mathcal{P}) \otimes \mathbb{Q} = \mathbb{Q}[f_\lambda(\alpha), \alpha^2]$.

Proof. At first let us proof that the polynomials $\{f_\lambda(\alpha)\}$, α^2 are algebraically independent.

Let $g \in \mathbb{Q}[x, y_1, y_2, \dots, y_s]$ be a polynomial such that $g(\alpha^2, f_1(\alpha), \dots, f_s(\alpha)) = 0$. We can write this equality as

$$g_0(f_1(\alpha), \dots, f_s(\alpha)) + g_2(f_1(\alpha), \dots, f_s(\alpha))\alpha^2 + \dots + g_{2t}(f_1(\alpha), \dots, f_s(\alpha))\alpha^{2t} = 0$$

for some integers s, t . Let us set $\alpha = 0$. Then we obtain

$$g_0(f_1, \dots, f_s) = 0$$

Since $\{f_\lambda\}$ are algebraically independent, $g_0 \equiv 0$. Since $\mathcal{D}^*[\alpha]$ is a polynomial ring, we can divide the equality by α^2 to obtain

$$g_2(f_1(\alpha), \dots, f_s(\alpha)) + g_4(f_1(\alpha), \dots, f_s(\alpha))\alpha^2 + \dots + g_{2t}(f_1(\alpha), \dots, f_s(\alpha))\alpha^{2t-2} = 0$$

Iterating this step in the end we obtain that all the polynomials $g_{2i} \equiv 0$, $i = 0, 1, \dots, 2t$, so $g \equiv 0$.

Now let us prove that the polynomials $\{f_\lambda(\alpha)\}$, α^2 generate $f(\mathcal{P}) \otimes \mathbb{Q}$.

Let $\psi(\alpha) = \psi_0 + \psi_1\alpha + \dots + \psi_n\alpha^n \in f(\mathcal{P}) \otimes \mathbb{Q}$ be a homogeneous element of degree $2n$. The functions $\{f_\lambda\}$ generate $\mathcal{D}^* \otimes \mathbb{Q}$, so we can find a homogeneous polynomial $g_0 \in \mathbb{Q}[y_1, \dots, y_s]$ such that

$$g_0(f_1, \dots, f_s) = \psi_0, \quad \deg f_i \leq 2n$$

Let us consider the element $\psi(\alpha) - g_0(f_1(\alpha), \dots, f_s(\alpha)) \in f(\mathcal{P}) \otimes \mathbb{Q}$. Since $\theta_1 = \theta_0 \frac{d}{2}$ for any

$$\theta(\alpha) = \theta_0 + \theta_1\alpha + \theta_2\alpha^2 + \dots + \theta_r\alpha^r \in f(\mathcal{P}) \otimes \mathbb{Q},$$

we obtain that

$$\psi(\alpha) - g_0(f_1(\alpha), \dots, f_s(\alpha)) = \hat{\psi}(\alpha)\alpha^2.$$

is a homogeneous element of degree $2n$. $\hat{\psi}(\alpha)\alpha^2 \in f(\mathcal{P}) \otimes \mathbb{Q}$, so $\hat{\psi}(-\alpha)\alpha^2\Phi(\alpha) = \hat{\psi}(\alpha)\alpha^2$. Since $\mathcal{D}^* \otimes \mathbb{Q}[\alpha]$ is a polynomial ring, $\hat{\psi}(-\alpha)\Phi(\alpha) = \hat{\psi}(\alpha)$, so $\hat{\psi}(\alpha) \in f(\mathcal{P}) \otimes \mathbb{Q}$ is a homogeneous element of degree $2(n-2)$.

Iterating this argument in the end we obtain the expression of $\psi(\alpha)$ as a polynomial in $\{f_\lambda(\alpha)\}$ and α^2 . □

This lemma proves the theorem. □

Let us note that dimension of the $(2n)$ -th graded component of the ring $f(\mathcal{P}) \otimes \mathbb{Q}$ is equal to the n -th Fibonacci number c_n .

Corollary 8.3. *There is an isomorphism of free polynomial algebras*

$$f(\mathcal{P}) \otimes \mathbb{Q} \simeq \mathbb{Q}[\text{LYN}_{\text{odd}}, \alpha^2] = \mathcal{N}_{\text{odd}} \otimes \mathbb{Q}[\alpha^2] \simeq \mathbb{Q}[\text{LYN}_{12}] = \mathcal{N}_{12} \otimes \mathbb{Q}.$$

Proof. Indeed, these three algebras are free polynomial algebras with the same dimensions of the corresponding graded components: dimension of the $(2n)$ -th graded component of each algebra is equal to c_n □

Let us denote by k_n the number of multiplicative generators of degree $2n$. According to Corollary 8.3 for $n \geq 3$ the number of generators k_n is equal to the number of Lyndon words of degree $2n$ in LYN_{odd} or in LYN_{12} .

For example, for small n we have

n	LYN_{12}^{2n}	$\text{LYN}_{\text{odd}}^{2n}$
3	[1,2]	[3]
4	[1,1,2]	[1,3]
5	[1,2,2], [1,1,1,2]	[5], [1,1,3]
6	[1,1,2,2], [1,1,1,1,2]	[1,5], [1,1,1,3]
7	[1,2,2,2], [1,1,2,1,2], [1,1,1,2,2], [1,1,1,1,1,2]	[7], [1,1,5], [1,3,3], [1,1,1,1,3],

Corollary 8.4. 1. $k_{n+1} \geq k_n$

2. $k_n \geq N_n - 2$, where N_n is the number of different decompositions of n into the sum of odd numbers.

Remark 8.5. It is a well-known fact, that the number of the decompositions of n into the sum of odd numbers is equal to the number of the decompositions of n into the sum of different numbers. For example,

$$\begin{array}{rclcl} 2 & = & 1 + 1 & = & 2 \\ 3 & = & 1 + 1 + 1, 3 & = & 1 + 2, 3 \\ 4 & = & 1 + 1 + 1 + 1, 1 + 3 & = & 1 + 3, 4 \\ 5 & = & 1 + 1 + 1 + 1 + 1, 1 + 1 + 3, 5 & = & 1 + 4, 2 + 3, 5 \end{array}$$

Proof. 1. $k_1 = k_2 = k_3 = 1$.

Let $n \geq 3$ and let $w = [a_1, \dots, a_k] \in \text{LYN}_{12}^{2n}$. Then $1w = [1, a_1, \dots, a_k] \in \text{LYN}_{12}^{2(n+1)}$. Indeed, for any proper tail $[a_i, \dots, a_k]$, $i \geq 1$ if $a_i > 1$, then $[a_i, \dots, a_k] > 1w$. If $a_i = 1$ then $i \neq k$ and $[a_{i+1}, \dots, a_k] > [a_1, \dots, a_k]$, since w is Lyndon. Thus $k_{n+1} \geq k_n$

2. Any decomposition $n = d_1 + \dots + d_k$ into the sum of odd numbers $d_1 \leq d_2 \leq \dots \leq d_k$ gives a Lyndon word $[d_1, \dots, d_k] \in \text{LYN}_{\text{odd}}$, except for the case, when $k > 1$, $d_1 = \dots = d_k = d \neq n$.

If $d \geq 5$, then the word $[1, d - 2, 1, d, \dots, d]$ is Lyndon. When $d = 1$ or $d = 3$ there can be no Lyndon word corresponding to the decomposition $n = 1 + \dots + 1$ or $n = 3 + \dots + 3$, as it happens for $n = 6$:

$$6 = 1 + 5 = 3 + 3 = 1 + 1 + 1 + 3 = 1 + 1 + 1 + 1 + 1 + 1$$

We have $k_6 = 2$ and $N_6 = 4$. Thus $k_n \geq N_n - 2$. □

Then we obtain the corollary:

Corollary 8.6. For any t , $|t| < \frac{\sqrt{5}-1}{2}$, there is the following decomposition of the generating series of Fibonacci numbers into the absolutely convergent infinite product:

$$\frac{1}{1-t-t^2} = 1 + t + t^2 + 2t^3 + 3t^4 + \dots = \sum_{n=0}^{\infty} c_n t^n = \prod_{i=1}^{\infty} \frac{1}{(1-t^i)^{k_i}},$$

The numbers k_n satisfy the properties $k_{n+1} \geq k_n \geq N_n - 2$, where N_n is the number of decompositions of n into the sum of odd summands.

Proof. Let $|t| < \frac{\sqrt{5}-1}{2}$. Then the series $\sum_{n=0}^{\infty} c_n t^n$ converges absolutely to $\frac{1}{1-t-t^2}$. Then

$$R_N = \sum_{n=N}^{\infty} c_n |t^n| \xrightarrow{N \rightarrow \infty} 0.$$

But we have:

$$S_n = \sum_{k=0}^{\infty} c_k t^k - \prod_{i=1}^{n-1} \frac{1}{(1-t^i)^{k_i}} = a_n t^n + a_{n+1} t^{n+1} + a_{n+2} t^{n+2} + \dots,$$

where $0 \leq a_i \leq c_i$. Then $|S_n| \leq R_n$, therefore $S_n \xrightarrow{n \rightarrow \infty} 0$, and the formula is valid.

The infinite product $\lim_{n \rightarrow \infty} \prod_{i=1}^n \frac{1}{(1-t^i)^{k_i}}$ converges absolutely if and only if the series $\sum_{n=1}^{\infty} |-k_n \ln(1-t^n)|$ converges. For $0 \leq t < \frac{\sqrt{5}-1}{2}$ this is true. In this case $0 \leq k_n t^n \leq -k_n \ln(1-t^n)$, therefore the power series $\sum_{n=1}^{\infty} k_n t^n$ converges and its radius of convergence is at least $\frac{\sqrt{5}-1}{2}$. Thus for $|t| < \frac{\sqrt{5}-1}{2}$ we have: $\sum_{n=1}^{\infty} k_n |t^n| < \infty$, and the equivalence $|-k_n \ln(1-t^n)| \sim k_n |t^n|$ implies the absolute convergence of the infinite product. □

Here are the examples in small dimensions:

n	generators	k_n	additive basis	c_n
1	[1]	1	[1]	1
2	[2]	1	[2], [1]*[1]	2
3	[1,2]	1	[1,2], [2]*[1], [1]*[1]*[1]	3
4	[1,1,2]	1	[1,1,2], [1,2]*[1], [2]*[2], [2]*[1]*[1], [1]*[1]*[1]*[1]	5
5	[1,2,2], [1,1,1,2]	2	[1,2,2], [1,1,1,2], [1,1,2]*[1], [1,2]*[2], [1,2]*[1]*[1], [2]*[2]*[1], [2]*[1]*[1]*[1], [1]*[1]*[1]*[1]*[1]	8
6	[1,1,2,2], [1,1,1,1,2]	2		13
7	[1,2,2,2], [1,1,2,1,2], [1,1,1,2,2], [1,1,1,1,1,2]	4		21

The corresponding products are:

$$\begin{aligned}
(1-t)(1-t^2) &= 1-t-t^2+t^3; \\
(1-t)(1-t^2)(1-t^3) &= 1-t-t^2+t^4+t^5-t^6; \\
(1-t)(1-t^2)(1-t^3)(1-t^4) &= 1-t-t^2+2t^5-t^8-t^9+t^{10}; \\
(1-t)(1-t^2)(1-t^3)(1-t^4)(1-t^5)^2 &= 1-t-t^2+2t^6+2t^7-t^8-t^9-2t^{10}- \\
&\quad -t^{11}-t^{12}+2t^{13}+2t^{14}-t^{18}-t^{19}+t^{20}.
\end{aligned}$$

The numbers k_i can be found in the following way:

$$\begin{aligned}
-\log(1-t-t^2) &= -\sum_{i=1}^{\infty} k_i \log(1-t^i); \\
\sum_{n=1}^{\infty} \frac{1}{n} \sum_{j=0}^n \binom{n}{j} t^{n+j} &= \sum_{i=1}^{\infty} k_i \sum_{r=1}^{\infty} \frac{t^{ri}}{r}
\end{aligned}$$

Thus for any N

$$\sum_{j=0}^{\lfloor \frac{N}{2} \rfloor} \frac{\binom{N-j}{j}}{N-j} = \frac{1}{N} \sum_{i|N} i k_i.$$

According to the Möbius inversion formula we obtain

$$N k_N = \sum_{d|N} \left(d \sum_{j=0}^{\lfloor \frac{d}{2} \rfloor} \frac{\binom{d-j}{j}}{d-j} \right) \cdot \mu\left(\frac{N}{d}\right),$$

where $\mu(n)$ is the Möbius function, that is

$$\mu(n) = \begin{cases} 1, & n = 1; \\ (-1)^r, & n = p_1 \dots p_r, \{p_i\} - \text{distinct prime numbers}; \\ 0, & n \text{ is not square-free.} \end{cases}$$

For example, if N is a prime number p , then

$$p k_p = -1 + p \sum_{j=0}^{\lfloor \frac{p}{2} \rfloor} \frac{\binom{p-j}{j}}{p-j}; \quad k_p = \sum_{j=1}^{\lfloor \frac{p}{2} \rfloor} \frac{\binom{p-j}{j}}{p-j}.$$

Each summand is an integer, since $\binom{p-j}{j} = \frac{(p-j-1)(p-j)}{j}$ is an integer, and $(p-j)$ and j are relatively prime numbers.

Thus $k_5 = 2$, $k_7 = 4$, $k_{11} = 18$, and so on.

9 Bayer-Billera Ring

Let us consider the free graded abelian group $BB \subset \mathcal{P}$, generated by 1 and all the polytopes $Q \in \Omega^n$, $n \geq 1$. Rank of the $(2n)$ -th graded component of this group is equal to c_n .

Since the determinant of the matrix K^n is equal to 1, the generalized f -polynomials $\{f(Q), Q \in \Omega^n\}$ form a basis of the $(2n)$ -th graded component of the ring $f(\mathcal{P})$.

So the composition of the inclusion $i : BB \subset \mathcal{P}$ and the mapping $f : \mathcal{P} \rightarrow \text{Qsym}[t_1, t_2, \dots][\alpha]$ is an isomorphism of the abelian groups BB and $f(\mathcal{P})$. This gives a projection $\pi : \mathcal{P} \rightarrow BB$

$$\pi(p) = x \in BB : f(p) = f(x).$$

It follows from the definition, that $\pi \circ i = 1$ on the space BB .

Theorem 9.1. *The projection $\pi : \mathcal{P} \rightarrow BB$ defined by the relation $f(\pi p) = f(p)$ gives the graded group BB the structure of a graded commutative associative ring with the multiplication $x \times_{BB} y = \pi(x \times y)$ such that $f(x \times_{BB} y) = f(x)f(y)$, and $BB \otimes \mathbb{Q}$ is a free polynomial algebra in a countable set of variables.*

Question 9.2. To describe the projection $\pi : \mathcal{P} \rightarrow BB$ in an efficient way.

Question 9.3. To find the multiplicative generators of $BB \otimes \mathbb{Q}$ in an efficient way.

Question 9.4. To describe the multiplication in BB in an efficient way.

10 Hopf comodule structures

Let R be a field or the ring $\mathbb{Z}$.

Definition 10.1. By a (left) *Hopf comodule* (or *Milnor comodule*) over a Hopf algebra X we mean an R -algebra M with a unit provided M is a comodule over X with a coaction $b : M \rightarrow X \otimes M$ such that $b(uv) = b(u)b(v)$, i.e. such that b is a homomorphism of rings.

10.1 Hopf comodule structure arising from the Hopf module structure

Lemma 10.2. *Let $\mathcal{H}$ be a graded connected torsionfree Hopf algebra over $\mathbb{Z}$ with finite rank of each graded component. If a ring $\mathcal{A}$ has the right graded Milnor module structure over $\mathcal{H}$, then $\mathcal{A}$ has a left graded Hopf comodule structure over the graded dual Hopf algebra $\mathcal{H}^*$ defined by the formula*

$$\Delta_{\mathcal{H}^*}(A) = \sum_{H_\omega \in \mathcal{H}} H_\omega^* \otimes AH_\omega$$

where $\{H_\omega\}$ is some basis in the graded group $\mathcal{H}$, and $\{H_\omega^*\}$ is the graded dual basis: $\langle H_\omega^*, H_\sigma \rangle = \delta_{\omega, \sigma}$.

Proof.

$$\begin{aligned} (1 \otimes \Delta_{\mathcal{H}^*}) \circ \Delta_{\mathcal{H}^*} A &= \sum_{\omega} H_\omega^* \otimes \left(\sum_{\sigma} H_\sigma^* \otimes (AH_\sigma) H_\omega \right) = \sum_{\omega, \sigma} H_\omega^* \otimes H_\sigma^* \otimes A \left(\sum_{\tau} \langle H_\tau^*, H_\omega H_\sigma \rangle H_\tau \right) = \\ &= \sum_{\tau} \left(\sum_{\omega, \sigma} H_\omega^* \otimes H_\sigma^* \langle H_\tau^*, H_\omega H_\sigma \rangle \right) \otimes AH_\tau = \sum_{\tau} \left(\sum_{\omega, \sigma} H_\omega^* \otimes H_\sigma^* \langle \Delta H_\tau^*, H_\omega \otimes H_\sigma \rangle \right) \otimes AH_\tau = \\ &= \sum_{\tau} \Delta H_\tau^* \otimes AH_\tau = (\Delta \otimes 1) \circ \Delta_{\mathcal{H}^*} A \end{aligned}$$

Since the augmentation in the ring $\mathcal{H}^*$ has the form $\varepsilon(\xi) = \xi(1)$, we see that

$$(\varepsilon \otimes 1) \Delta_{\mathcal{H}^*} A = (\varepsilon \otimes 1) \sum_{\omega} H_\omega^* \otimes AH_\omega = \sum_{\omega} H_\omega^*(1) \otimes AH_\omega = H_{\omega_0}^*(1) \otimes AH_{\omega_0} = 1 \otimes A,$$

where H_{ω_0} is a basis in $\mathcal{H}^0$.

Therefore $\Delta_{\mathcal{H}^*}$ is a comodule structure.

Now

$$\begin{aligned}
\Delta_{\mathcal{H}^*}(AB) &= \sum_{\omega} H_{\omega}^* \otimes (AB)H_{\omega} = \sum_{\omega} H_{\omega}^* \otimes \left(\sum_i (AH'_{\omega,i})(BH''_{\omega,i}) \right) = \\
&= \sum_{\omega} H_{\omega}^* \otimes \left(\sum_i \left(A \sum_{\sigma} \langle H_{\sigma}^*, H'_{\omega,i} \rangle H_{\sigma} \right) \left(B \sum_{\tau} \langle H_{\tau}^*, H''_{\omega,i} \rangle H_{\tau} \right) \right) = \\
&= \sum_{\sigma, \tau} \left(\sum_{\omega} \left(\sum_i \langle H_{\sigma}^*, H'_{\omega,i} \rangle \langle H_{\tau}^*, H''_{\omega,i} \rangle \right) H_{\omega}^* \right) \otimes (AH_{\sigma})(BH_{\tau}) = \\
&= \sum_{\sigma, \tau} \left(\sum_{\omega} \langle H_{\sigma}^* \cdot H_{\tau}^*, H_{\omega} \rangle H_{\omega}^* \right) \otimes (AH_{\sigma})(BH_{\tau}) = \sum_{\sigma, \tau} H_{\sigma}^* \cdot H_{\tau}^* \otimes (AH_{\sigma})(BH_{\tau}) = (\Delta_{\mathcal{H}^*}A) \cdot (\Delta_{\mathcal{H}^*}B),
\end{aligned}$$

where $\Delta H_{\omega} = \sum_i H'_{\omega,i} \otimes H''_{\omega,i}$. This equality finishes the proof. $\square$

Remark 10.3. This Hopf comodule structure does not depend on the choice of a graded basis in the graded group $\mathcal{H}$, since we can write it as

$$\Delta_{\mathcal{H}^*}A = (1 \otimes A) \left(\sum_{\omega} H_{\omega}^* \otimes H_{\omega} \right),$$

and $\sum_{\omega} H_{\omega}^* \otimes H_{\omega}$ defines in each graded component $\mathcal{H}^n$ the identity operator in $(\mathcal{H}^n)^* \otimes \mathcal{H}^n \simeq \text{Hom}_{\mathbb{Z}}(\mathcal{H}^n, \mathcal{H}^n)$.

Remark 10.4. The similar argument proves that if a ring $\mathcal{A}$ has the left graded Milnor module structure over $\mathcal{H}$, then A has a right graded Hopf comodule structure over the graded dual Hopf algebra $\mathcal{H}^*$ defined by the formula

$$\Delta_{\mathcal{H}^*}(A) = \sum_{H_{\omega} \in \mathcal{H}} H_{\omega} A \otimes H_{\omega}^*.$$

Corollary 10.5. *There is a natural left Hopf comodule structure on the ring $\mathbf{R}$ over the Hopf algebra $\mathcal{Q}\text{sym}$ induced by the right Hopf module structure over the Hopf algebra $\mathcal{Z}$:*

$$\Delta_{\mathcal{R}}(P) = \sum_{\omega} M_{\omega} \otimes PZ_{\omega} = \sum_{k \geq 0} \sum_{(j_1, \dots, j_k)} M_{(j_1, \dots, j_k)} \otimes (PZ_{j_1} \dots Z_{j_k}) = \sum_{k \geq 0} \sum_{(j_1, \dots, j_k)} M_{(j_k, \dots, j_1)} \otimes (d_{j_1} \dots d_{j_k} P)$$

There are natural right Hopf comodule structures $\Delta_{\mathcal{L}}$ and $\Delta_{\mathcal{D}^*}$ on the ring $\mathbf{R}$ over the Hopf algebras $\mathcal{Q}\text{sym}$ and $\mathcal{D}^*$ induced by the left Hopf module structures over the Hopf algebras $\mathcal{Z}$ and $\mathcal{D}$.

These structures are compatible in the following sense:

$$\Delta_{\mathcal{L}} = \tau \circ (* \otimes 1) \circ \Delta_{\mathcal{R}} \tag{20}$$

$$(1 \otimes \mathcal{L}^*) \circ \Delta_{\mathcal{D}^*} = \Delta_{\mathcal{L}} \tag{21}$$

where $\tau: \mathcal{Q}\text{sym} \otimes \mathbf{R} \rightarrow \mathbf{R} \otimes \mathcal{Q}\text{sym}$ is a homomorphism that interchanges the tensor factors

$$\tau(M_{\omega} \otimes P) = P \otimes M_{\omega}.$$

We have

$$\Delta_{\mathcal{L}}P = \sum_{k \geq 0} \sum_{(j_1, \dots, j_k)} (d_{j_1} \dots d_{j_k} P) \otimes M_{(j_1, \dots, j_k)}$$

Proof. The existence of these structures and the explicit formulas follow immediately from Lemma 10.2 and Corollary 10.5. The first relation follows easily from the formulas for $\Delta_{\mathcal{L}}$ and $\Delta_{\mathcal{R}}$.

Since $\mathcal{D} \simeq \mathcal{Z}/J_{\mathcal{U}}$ is a free abelian group, we have $\mathcal{Z} = \mathcal{Z}' \oplus J_{\mathcal{U}}$. Let us choose graded bases in both subgroups: $\{H_{\beta}\}$ in $\mathcal{Z}'$, and $\{H_{\gamma}\}$ in $J_{\mathcal{U}}$. Then $\{H_{\beta}, H_{\gamma}\}$ form a graded basis in $\mathcal{Z}$.

Let us note that $\mathcal{L}^*(\mathcal{D}^*) = \{\psi \in \mathcal{Q}\text{sym} : \psi|_{J_{\mathcal{U}}} = 0\}$.

Then the elements of the dual basis $\{H_\beta^*\}, \{H_\gamma^*\}$ form a basis in $\mathcal{Qsym}$, while the functions $\{H_\beta^*\}$ form a basis in $\mathcal{L}^*(\mathcal{D}^*)$. Let $H_\beta^* = \mathcal{L}^* D_\beta^*$. Then the operators $\{D_\beta = \mathcal{L} H_\beta\}$ form a graded basis in $\mathcal{D}$, dual to the basis $\{D_\beta^*\}$.

Now for any $P \in \mathcal{R}$ we have

$$\Delta_{\mathcal{L}} P = \sum_{\beta} (H_\beta P) \otimes H_\beta^* + \sum_{\gamma} (H_\gamma P) \otimes H_\gamma^*$$

But $H_\gamma P = 0$ for all γ , therefore,

$$\Delta_{\mathcal{L}} P = \sum_{\beta} H_\beta P \otimes H_\beta^* = \sum_{\beta} (\mathcal{L} H_\beta) P \otimes H_\beta^* = \sum_{\beta} D_\beta P \otimes \mathcal{L}^* D_\beta^* = (1 \otimes \mathcal{L}^*) \circ \Delta_{\mathcal{D}^*} P$$

□

We have the series of mappings

$$\Phi(t_1): \mathcal{R} \rightarrow \mathcal{R}[t_1], \Phi_2(t_1, t_2) = \Phi(t_2)\Phi(t_1): \mathcal{R} \rightarrow \mathcal{Qsym}[t_1, t_2] \otimes \mathcal{R},$$

...

$$\Phi_n(t_1, \dots, t_n) = \Phi(t_n)\Phi(t_{n-1}) \dots \Phi(t_1): \mathcal{R} \rightarrow \mathcal{Qsym}[t_1, \dots, t_n] \otimes \mathcal{R}$$

Taking the inverse limit we obtain the mapping

$$\Phi_\infty: \mathcal{R} \rightarrow \mathcal{Qsym} \otimes \mathcal{R},$$

such that for any element $P \in \mathcal{R}$ of graduation $2n$

$$\Phi_\infty P = \sum_{k=0}^n \sum_{(j_1, \dots, j_k)} M_{(j_k, \dots, j_1)} \otimes (d_{j_1} \dots d_{j_k} P)$$

Thus we have proved the following fact.

Proposition 10.6.

$$\Phi_\infty = \Delta_{\mathcal{R}}$$

10.2 Ring homomorphisms

Definition 10.7. Any ring homomorphism $h: \mathcal{R} \rightarrow \mathcal{A}$ induces the ring homomorphism

$$\Phi_h: \mathcal{R} \rightarrow \mathcal{Qsym} \otimes \mathcal{A}, \Phi_h = (1 \otimes h) \circ \Delta_{\mathcal{R}}$$

Let us denote by $\Phi_{h,r}$ the homomorphism $\mathcal{R} \rightarrow \mathcal{Qsym}[t_1, \dots, t_r] \otimes \mathcal{A}$.

$$\Phi_{h,r}(P) = \Phi_h(P)(t_1, \dots, t_r, 0, 0, \dots)$$

Example 10.8. Let us consider several examples.

1. $\xi_\alpha: \mathcal{P} \rightarrow \mathbb{Z}[\alpha]$ induces the homomorphism $\Phi_{\xi_\alpha}: \mathcal{P} \rightarrow \mathcal{Qsym}[\alpha]$. This homomorphism coincides with the generalized f -polynomial.
2. $\varepsilon_\alpha: \mathcal{RP} \rightarrow \mathbb{Z}[\alpha]$ induces the ring homomorphism $\Phi_{\varepsilon_\alpha}: \mathcal{RP} \rightarrow \mathcal{Qsym}[\alpha]$. Let us denote this ring homomorphism by $f_{\mathcal{RP}}$.
3. $\varepsilon_0: \mathcal{RP} \rightarrow \mathbb{Z}$ induces the homomorphism $\Phi_{\varepsilon_0}: \mathcal{RP} \rightarrow \mathcal{Qsym}$. Then $\Phi_{\varepsilon_0} = F^*$, where F is the Ehrenborg F -quasi-symmetric function introduced in [Ehr]:

$$F(P) = \sum_{\hat{0}=x_0 < x_1 < \dots < x_{k+1}=\hat{1}} M_{(\rho(x_0, x_1), \rho(x_1, x_2), \dots, \rho(x_k, x_{k+1}))} = \sum_{0 \leq a_1 < \dots < a_k \leq n-1} f_{a_1, \dots, a_k}(P) M_{(a_1+1, a_2-a_1, \dots, n-a_k)}$$

where P is an n -dimensional polytope, the sum ranges over all chains from $\hat{0} = \emptyset$ to $\hat{1} = P$ in the face lattice $L(P)$, and $f_{a_1, \dots, a_k}$ are flag numbers.

In fact, F is defined on the Rota-Hopf algebra $\mathcal{R}$ of graded posets, and it was proved in [Ehr] that it is a Hopf algebra homomorphism $\mathcal{R} \rightarrow \mathcal{Qsym}$ such that $F(P^*) = F(P)^*$.

Proposition 10.9. *For any polytope $P \in \mathcal{P}$ we have*

$$f_{\mathcal{RP}}(P) = F^*(P) + \alpha f(P)$$

Proof. Indeed, let P be an n -dimensional polytope. Then

$$f_{\mathcal{RP}}(P) = \sum_{k=0}^n \sum_{0 \leq a_1 < \dots < a_k \leq n-1} f_{a_1, \dots, a_k} (M_{(n-a_k, \dots, a_1+1)} + \alpha^{a_1+1} M_{(n-a_k, \dots, a_2-a_1)})$$

□

Corollary 10.10. *For any two polytopes $P, Q \in \mathcal{P}$ we have*

$$f(P * Q) = f(P) \cdot F^*(Q) + F^*(P) \cdot f(Q) + \alpha f(P) \cdot f(Q)$$

Proof. Since

$$f_{\mathcal{RP}}(P * Q) = f_{\mathcal{RP}}(P) \cdot f_{\mathcal{RP}}(Q), \quad \text{and} \quad F^*(P * Q) = F^*(P) \cdot F^*(Q),$$

we obtain

$$F^*(P) \cdot F^*(Q) + \alpha f(P * Q) = (F^*(P) + \alpha f(P)) \cdot (F^*(Q) + \alpha f(Q))$$

Removing the brackets, canceling the equal summands, and dividing by α , we obtain the required formula. □

Corollary 10.11.

$$f(CP) = F^*(P) + (\alpha + \sigma_1)f(P)$$

Theorem 10.12. • *For $r \geq n$ the image $f_r(\mathcal{P}^{2n}) \subset \text{Qsym}[t_1, \dots, t_r][\alpha]$ consists of homogeneous polynomials g of degree $2n$ satisfying the equations*

1.

$$\begin{aligned} g(\alpha, t_1, -t_1, t_3, \dots, t_r) &= g(\alpha, 0, 0, t_3, \dots, t_r) \\ g(\alpha, t_1, t_2, -t_2, \dots, t_r) &= g(\alpha, t_1, 0, 0, \dots, t_r) \\ &\dots \\ g(\alpha, t_1, \dots, t_{r-1}, -t_{r-1}) &= g(\alpha, t_1, \dots, 0, 0) \end{aligned} \tag{22}$$

2.

$$g(-\alpha, t_1, t_2, \dots, t_{r-1}, \alpha) = g(\alpha, t_1, t_2, \dots, t_{r-1}, 0) \tag{23}$$

- *For $r \geq n$ the image $f_{\mathcal{RP}, r}(\mathcal{RP}^{2n}) \subset \text{Qsym}[t_1, \dots, t_r][\alpha]$ consists of homogeneous polynomials g of degree $2n$ satisfying the following conditions:*

1. *The equations of type (22)*

2. *The equation*

$$g(-\alpha, t_1, t_2, \dots, t_{r-1}, \alpha) = g(0, t_1, t_2, \dots, t_{r-1}, 0)$$

- *For $r \geq n$ the image $F_r(\mathcal{RP}^{2n}) \subset \text{Qsym}[t_1, \dots, t_r]$ consists of homogeneous polynomials g of degree $2n$ satisfying the equations of type (22).*

In all the cases the equations are equivalent to the Bayer-Billera (generalized Dehn-Sommerville) relations [BB]

Proof. The first statement of the theorem is exactly Theorem 6.3.

For the second and the third statements we need a lemma corresponding to Lemma 6.4

Lemma 10.13. *Let P be an n -dimensional polytope. Then*

1. The equation

$$f_{\mathcal{RP},r}(\alpha, t_1, \dots, t_q, -t_q, \dots, t_r) = f_{\mathcal{RP},r}(\alpha, t_1, \dots, 0, 0, \dots, t_r)$$

is equivalent to the generalized Dehn-Sommerville relations

$$\sum_{j=a_t+1}^{a_{t+1}-1} (-1)^{j-a_t-1} f_{a_1, \dots, a_t, j, a_{t+1}, \dots, a_k} = (1 + (-1)^{a_{t+1}-a_t}) f_{a_1, \dots, a_t, a_{t+1}, \dots, a_k}$$

for $-1 \leq a_1 < a_2 < \dots < a_k < n$, and $1 \leq k \leq \min\{r-1, n\}$, $k+1-q \leq t \leq r-q$.

2. The equation

$$f_{\mathcal{RP},r}(-\alpha, t_1, \dots, t_{r-1}, \alpha) = f_{\mathcal{RP},r}(0, t_1, \dots, t_{r-1}, 0)$$

is equivalent to the generalized Dehn-Sommerville relations

$$\sum_{j=0}^{a_1-1} (-1)^j f_{j, a_1, \dots, a_k} = (1 + (-1)^{a_1-1}) f_{a_1, \dots, a_k}$$

for $0 \leq k \leq \min\{r-1, n-1\}$.

3. The equation

$$F_r^*(t_1, \dots, t_q, -t_q, \dots, t_r) = F_r^*(t_1, \dots, 0, 0, \dots, t_r)$$

is equivalent to the generalized Dehn-Sommerville relations

$$\sum_{j=a_t+1}^{a_{t+1}-1} (-1)^{j-a_t-1} f_{a_1, \dots, a_t, j, a_{t+1}, \dots, a_k} = (1 + (-1)^{a_{t+1}-a_t}) f_{a_1, \dots, a_t, a_{t+1}, \dots, a_k}$$

for $0 \leq a_1 < \dots < a_k$, and $0 \leq k \leq \min\{r-2, n-1\}$, $k+2-q \leq t \leq r-q$.

Proof. Items 1. and 3. are proved in the same manner as Lemma 6.4. Let us consider the relation of item 2.: $f_{\mathcal{RP},r}(-\alpha, t_1, \dots, t_{r-1}, \alpha) = f_{\mathcal{RP},r}(0, t_1, \dots, t_{r-1}, 0)$:

$$\begin{aligned} &= (-\alpha)^{n+1} + \sum_{k=1}^{\min\{r-1, n+1\}} \sum_{-1 \leq a_1 < \dots < a_k \leq n-1} f_{a_1, \dots, a_k} (-\alpha)^{a_1+1} \left(\sum_{1 \leq l_1 < \dots < l_k \leq r-1} t_{l_1}^{n-a_k} \dots t_{l_k}^{a_2-a_1} \right) + \\ &+ \sum_{k=1}^{\min\{r, n+1\}} \sum_{-1 \leq a_1 < \dots < a_k \leq n-1} f_{a_1, \dots, a_k} (-\alpha)^{a_1+1} \left(\sum_{1 \leq l_1 < \dots < l_k = r} t_{l_1}^{n-a_k} \dots t_{l_{k-1}}^{a_3-a_2} \alpha^{a_2-a_1} \right) = \\ &\sum_{k=1}^{\min\{r-1, n+1\}} \sum_{-1=a_1 < \dots < a_k \leq n-1} f_{a_1, \dots, a_k} \left(\sum_{1 \leq l_1 < \dots < l_k \leq r-1} t_{l_1}^{n-a_k} \dots t_{l_k}^{a_2-a_1} \right) \end{aligned}$$

This is equivalent to the relations

$$f_{a_1, \dots, a_k} + \sum_{j=0}^{a_1-1} (-1)^{j+1} f_{j, a_1, \dots, a_k} + (-1)^{a_1+1} f_{a_1, \dots, a_k} = 0$$

for $0 \leq a_1 < \dots < a_k$, and $0 \leq k \leq \min\{r-1, n-1\}$. The case $k=0$ again corresponds to the Euler formula

$$1 - f_0 + f_1 + \dots + (-1)^n f_{n-1} + (-1)^{n+1} = 0.$$

□

Remark 10.14. Let us mention that in the case of $f_{\mathcal{RP},r}$ and F_r^* the equations of type (22) contain all the generalized Dehn-Sommerville relations, and these equations for $f_{\mathcal{RP},r}$ imply the equation (23). This follows from the fact that a_1 can be equal to -1 , which is not possible in the case of the generalized f -polynomial.

Now the end of the proof of the theorem is the same as for Theorem 6.3. $\square$

Proposition 10.15. *The correspondence $f_{\mathcal{RP}} \rightarrow f_{\mathcal{RP}}|_{\alpha=0} = F^*$ is an isomorphism of the images $f_{\mathcal{RP}}(\mathcal{RP})$ and $F^*(\mathcal{RP})$.*

Proof. Indeed, it is clear that this mapping is an epimorphism. On the other hand, let $F^*(P) = 0$. Then all the flag numbers of the element $P \in \mathcal{RP}$ are equal to 0. Therefore, $f_{\mathcal{RP}}(P) = 0$, so the mapping is a monomorphism. $\square$

Corollary 10.16. 1. *The ring homomorphism $F^* : \mathcal{RP} \rightarrow \mathcal{R}^*(\mathcal{D}^*) \subset \mathcal{Qsym}$ is an epimorphism,*

2. *$F^*(\mathcal{RP}) \otimes \mathbb{Q}$ is a free polynomial algebra with dimension of the $(2n)$ -th graded component equal to the $(n-1)$ -th Fibonacci number c_{n-1} ($c_0 = c_1 = 1$, $c_{n+1} = c_n + c_{n-1}$, $n \geq 1$).*

Proof. 1. Let $g \in \mathcal{R}^*(\mathcal{D}^*)$ be a quasi-symmetric function of degree $2n$. Then according to Theorem 7.3 $g_n(t_1, \dots, t_n) = g(t_1, \dots, t_n, 0, 0, \dots)$ satisfies the relations of type (22). Therefore $g_n(t_1, \dots, t_n) = F_n^*(P)$ for some $P \in \mathcal{RP}^{2n}$. Since the mapping $\mathcal{Qsym} \rightarrow \mathcal{Qsym}[t_1, \dots, t_n], t_j \rightarrow 0, j > n$ is injective on the graduation $(2n)$, we see that $g = F^*(P)$.

2. We see that $F^*(\mathcal{RP}) = \mathcal{R}^*(\mathcal{D}^*)$. Therefore,

$$F^*(\mathcal{RP}) \otimes \mathbb{Q} \simeq \mathcal{D}^* \otimes \mathbb{Q} \simeq \mathbb{Q}[\text{LYN}_{\text{odd}}],$$

and $\dim F^*(\mathcal{RP}^{2n}) \otimes \mathbb{Q} = c_{n-1}$ according to Proposition 7.7. $\square$

Remark 10.17. Let us mention that $\deg L(P^n) = 2(n+1)$, therefore, $\text{rank } F^*(\mathcal{RP}^{2(n+1)}) = c_n = \text{rank } f(\mathcal{P}^{2n})$

10.3 Natural Hopf comodule structure

$\mathcal{P}$ is not a Hopf algebra, but it turns out that there is a natural Hopf comodule structure on the ring $\mathcal{P}$ over the Hopf algebra $\mathcal{RP}$.

Proposition 10.18. *Formula*

$$\Delta_{\mathcal{RP}} P = \sum_{F \subseteq P} F \otimes (P/F)$$

defines a natural graded right Hopf comodule structure $\Delta_{\mathcal{RP}} : \mathcal{P} \rightarrow \mathcal{P} \otimes \mathcal{RP}$ on the ring $\mathcal{P}$ over the Hopf algebra $\mathcal{RP}$.

Proof. We should proof that the following two diagrams commute

$$\begin{array}{ccc} \mathcal{P} & \xrightarrow{\Delta_{\mathcal{RP}}} & \mathcal{P} \otimes \mathcal{RP} \\ \downarrow \Delta_{\mathcal{RP}} & & \downarrow \text{id} \otimes \Delta \\ \mathcal{P} \otimes \mathcal{RP} & \xrightarrow{\Delta_{\mathcal{RP}} \otimes \text{id}} & \mathcal{P} \otimes \mathcal{RP} \otimes \mathcal{RP} \end{array} \quad \begin{array}{ccc} \mathcal{P} & & \\ \downarrow \Delta_{\mathcal{RP}} & \searrow \text{id} \otimes 1 & \\ \mathcal{P} \otimes \mathcal{RP} & \xrightarrow{\text{id} \otimes \varepsilon} & \mathcal{P} \otimes \mathbb{Z} \end{array}$$

and that $\Delta_{\mathcal{RP}}$ is a ring homomorphism. Indeed, we have

$$\begin{aligned} (\Delta_{\mathcal{RP}} \otimes \text{id}) \circ \Delta_{\mathcal{RP}} P &= (\Delta_{\mathcal{RP}} \otimes \text{id}) \left(\sum_{F \subseteq P} F \otimes (P/F) \right) = \sum_{F \subseteq P} \left(\sum_{G \subseteq F} G \otimes (F/G) \right) \otimes P/F = \\ &= \sum_{G \subseteq P} G \otimes \left(\sum_{G \subseteq F \subseteq P} F/G \otimes P/F \right) = \sum_{G \subseteq P} G \otimes \Delta(P/G) = (\text{id} \otimes \Delta) \circ \Delta_{\mathcal{RP}} P, \end{aligned}$$

and

$$(\text{id} \otimes \varepsilon) \circ \Delta_{\mathcal{RP}}(P) = \sum_{F \subseteq P} F \otimes \varepsilon(P/F) = P \otimes 1,$$

Let us check that $\Delta_{\mathcal{RP}}$ is a homomorphism of rings $\mathcal{P} \rightarrow \mathcal{P} \otimes \mathcal{RP}$:

$$\begin{aligned}\Delta_{\mathcal{RP}}(P \times Q) &= \sum_{F \times G \subseteq P \times Q} (F \times G) \otimes (P \times Q / F \times G) = \sum_{F \subseteq P, G \subseteq Q} (F \times G) \otimes (P/F * Q/G) = \\ &= \left(\sum_{F \subseteq P} F \otimes P/F \right) \cdot \left(\sum_{G \subseteq Q} G \otimes Q/G \right) = \Delta_{\mathcal{RP}}(P) \cdot \Delta_{\mathcal{RP}}(Q) \\ \Delta_{\mathcal{RP}}(\text{pt}) &= \text{pt} \otimes (\text{pt} / \text{pt}) = \text{pt} \otimes \emptyset = 1 \otimes 1.\end{aligned}$$

□

Corollary 10.19. • Any ring homomorphism $\mathcal{P} \rightarrow \mathcal{A}$, where $\mathcal{A}$ is a ring, induces the ring homomorphism $\mathcal{P} \rightarrow \mathcal{A} \otimes \mathcal{RP}$

- Any linear homomorphism $\psi: \mathcal{RP} \rightarrow \mathbb{Z}$ induces the linear operator $\mathcal{P} \rightarrow \mathcal{P}$, which is a ring homomorphism, if ψ is a ring homomorphism.

Example 10.20. The homomorphism $\xi_\alpha: \mathcal{P} \rightarrow \mathbb{Z}[\alpha]$ defines the ring homomorphism $l_\alpha: \mathcal{P} \rightarrow \mathcal{RP}[\alpha]$:

$$l_\alpha(P^n) = \sum_{F \subseteq P} \alpha^{\dim F} P/F.$$

Let us recall that in the case of a simple n -polytope a face polytope of an i -dimensional face is a simplex $\Delta^{n-i-1} = x^{n-i}$. Therefore,

$$l_\alpha(P^n) = \sum_{i=0}^n f_i \alpha^i x^{n-i} = f_1(\alpha, x).$$

This is a homogeneous f -polynomial in two variables defined in [Buch].

Proposition 10.21.

$$\Delta_{\mathcal{RP}}(P * Q) = (1 \otimes P) \cdot \Delta_{\mathcal{RP}}(Q) + \Delta_{\mathcal{RP}}(P) \cdot (1 \otimes Q) + \Delta_{\mathcal{RP}}(P)(* \otimes *) \Delta_{\mathcal{RP}}(Q) \quad (24)$$

Proof.

$$\begin{aligned}\Delta_{\mathcal{RP}}(P * Q) &= \sum_{G \subseteq Q} (\emptyset * G) \otimes (P * Q / \emptyset * G) + \sum_{F \subseteq P} (F * \emptyset) \otimes (P * Q / F * \emptyset) + \\ &+ \sum_{F \subseteq P, G \subseteq Q} (F * G) \otimes (P * Q / F * G) = \sum_{G \subseteq Q} G \otimes (P * (Q/G)) + \sum_{F \subseteq P} F \otimes ((P/F) * Q) + \\ &+ \sum_{F \subseteq P, G \subseteq Q} (F * G) \otimes (P/F * Q/G) = (1 \otimes P) \Delta_{\mathcal{RP}}(Q) + \Delta_{\mathcal{RP}}(P)(1 \otimes Q) + \Delta_{\mathcal{RP}}(P)(* \otimes *) \Delta_{\mathcal{RP}}(Q)\end{aligned}$$

□

Now if we apply $\xi_\alpha \otimes 1$ to the formula (24), then we obtain the corollary:

Corollary 10.22.

$$l_\alpha(P * Q) = P * l_\alpha(Q) + l_\alpha(P) * Q + \alpha l_\alpha(P) \cdot l_\alpha(Q)$$

10.4 Interrelation

Let us remind that the left action of the Leibnitz-Hopf algebra $\mathcal{Z}$ in the ring $\mathcal{R}$ induces the right Hopf comodule structure

$$\Delta_{\mathcal{L}} P^n = \sum_{k=0}^n \sum_{(j_1, \dots, j_k)} (d_{j_1} \dots d_{j_k} P) \otimes M_{(j_1, \dots, j_k)},$$

Proposition 10.23. *The following diagram commutes:*

$$\begin{array}{ccc} \mathcal{P} & \xrightarrow{\Delta_{\mathcal{RP}}} & \mathcal{P} \otimes \mathcal{RP} \\ & \searrow \Delta_{\mathcal{L}} & \downarrow 1 \otimes F \\ & & \mathcal{P} \otimes \mathcal{Qsym} \end{array}$$

Proof. Let P be an n -dimensional polytope. Then

$$\begin{aligned} \Delta_{\mathcal{L}}(P) &= \sum_{k=0}^n \sum_{(j_1, \dots, j_k)} (d_{j_1} \dots d_{j_k} P) \otimes M_{(j_1, \dots, j_k)} = \\ &= \sum_{k=0}^n \sum_{(j_1, \dots, j_k)} \sum_{F^{n-(j_1+\dots+j_k)}} \left(F^{n-(j_1+\dots+j_k)} \sum_{F^{n-(j_1+\dots+j_k)} \subset F^{n-(j_2+\dots+j_k)} \subset \dots \subset F^{n-j_k} \subset P} 1 \right) \otimes M_{(j_1, \dots, j_k)} = \\ &= \sum_{r=0}^n \sum_{F^r \subseteq P} F^r \otimes \left(\sum_{k=0}^{n-r} \sum_{(j_1, \dots, j_k): j_1+\dots+j_k=n-r} \sum_{F^r \subset F^{r+j_1} \subset \dots \subset F^{n-j_k} \subset P} M_{(j_1, \dots, j_k)} \right) = \\ &= \sum_{F \subseteq P} F \otimes F(P/F) = (1 \otimes F) \circ \Delta_{\mathcal{RP}}(P) \end{aligned}$$

□

Corollary 10.24. *For any polytope $P \in \mathcal{P}$ we have*

$$F^*(l_\alpha P) = f(P)$$

Proof. We have $(\xi_\alpha \otimes 1) \circ \Delta_{\mathcal{L}} P = f(P)^*$. So if we apply the homomorphism $\xi_\alpha \otimes 1$ to the diagram above, we obtain $F \circ l_\alpha = (\xi_\alpha \otimes 1) \circ \Delta_{\mathcal{L}} = f^*$. Therefore, $F^* \circ l_\alpha = f$. □

Corollary 10.25. *The following diagram commutes:*

$$\begin{array}{ccc} \mathcal{P} & \xrightarrow{f^*} & \mathcal{Qsym}[\alpha] \\ \Delta_{\mathcal{RP}} \downarrow & & \downarrow \Delta \\ \mathcal{P} \otimes \mathcal{RP} & \xrightarrow{f^* \otimes F} & \mathcal{Qsym}[\alpha] \otimes \mathcal{Qsym} \end{array}$$

Proof. 1. Consider the commutative diagram

$$\begin{array}{ccccc} \mathcal{P} & \xrightarrow{\Delta_{\mathcal{RP}}} & \mathcal{P} \otimes \mathcal{RP} & \xrightarrow{\Delta_{\mathcal{L}} \otimes 1} & P \otimes \mathcal{Qsym} \otimes \mathcal{RP} \\ & \searrow \Delta_{\mathcal{L}} & \downarrow 1 \otimes F & & \downarrow 1 \otimes 1 \otimes F \\ & & \mathcal{P} \otimes \mathcal{Qsym} & \xrightarrow{\Delta_{\mathcal{L}} \otimes 1} & P \otimes \mathcal{Qsym} \otimes \mathcal{Qsym} \\ & & & \searrow & \downarrow \xi_\alpha \otimes 1 \otimes 1 \\ & & & & \mathbb{Z}[\alpha] \otimes \mathcal{Qsym} \otimes \mathcal{Qsym} \end{array}$$

Then

$$(\xi_\alpha \otimes 1 \otimes 1) \circ (\Delta_{\mathcal{L}} \otimes 1) \circ \Delta_{\mathcal{L}} = (\xi_\alpha \otimes 1 \otimes 1) \circ (1 \otimes \Delta) \circ \Delta_{\mathcal{L}} = \Delta \circ f^*$$

while

$$(\xi_\alpha \otimes 1 \otimes 1) \circ (1 \otimes 1 \otimes F) \circ (\Delta_{\mathcal{L}} \otimes 1) \circ \Delta_{\mathcal{RP}} = (f^* \otimes F) \circ \Delta_{\mathcal{RP}}$$

Since these two compositions correspond to two pathes in the diagram, they are equal.

2. Let us consider also a direct proof. We know that

$$\Delta(M_{(b_1, \dots, b_k)}) = 1 \otimes M_{(b_1, \dots, b_k)} + M_{(b_1)} \otimes M_{(b_2, \dots, b_k)} + \dots + M_{(b_1, \dots, b_{k-1})} \otimes M_{(b_k)} + M_{(b_1, \dots, b_k)} \otimes 1$$

and

$$f(\alpha, t_1, t_2, \dots)(P^n)^* = \alpha^n + \sum_{F^{a_1} \subset \dots \subset F^{a_k}} \alpha^{a_1} M_{(a_2 - a_1, \dots, n - a_k)} = \sum_{F^{a_1} \subset \dots \subset F^{a_k} \subset P^n} \alpha^{a_1} M_{(a_2 - a_1, \dots, n - a_k)}$$

Then

$$\begin{aligned} \Delta(f(P^n)^*) &= \sum_{F^{a_1} \subset \dots \subset G^{a_s} \subset \dots \subset F^{a_k} \subset P^n} \alpha^{a_1} M_{(a_2 - a_1, \dots, a_s - a_{s-1})} \otimes M_{(a_{s+1} - a_s, \dots, n - a_k)} = \\ &= \sum_{G^{a_s} \subseteq P^n} \left(\sum_{F^{a_1} \subset \dots \subset F^{a_{s-1}} \subset G^{a_s}} \alpha^{a_1} M_{(a_2 - a_1, \dots, a_s - a_{s-1})} \right) \otimes \left(\sum_{G^{a_s} \subset F^{a_{s+1}} \subset \dots \subset F^{a_k} \subset P^n} M_{(a_{s+1} - a_s, \dots, n - a_k)} \right) = \\ &= \sum_{G \subseteq P^n} f^*(G) \otimes F(P^n/G) = (f^* \otimes F) \circ \Delta_{\mathcal{RP}}(P^n). \end{aligned}$$

□

11 Operators B and C

We know that the *cone* and the *bipyramid* operations can be considered as linear operators on the rings $\mathcal{P}$ and $\mathcal{RP}$. It turns out that there are corresponding operations on the rings $\mathcal{Qsym}[\alpha]$ and $\mathcal{Qsym}$, which commute with the homomorphisms $f, f_{\mathcal{RP}}$, and F^* .

11.1 The case of $\mathcal{P}$

At first let us consider the ring $\mathcal{P}$. We have

$$[C, d] = 1 + \xi_0$$

For $k \geq 2$ we have the similar situation.

$$d_k C P^n = \begin{cases} C d_k P^n + d_{k-1} P^n, & k < n+1, \\ 1 + d_{k-1} P^n = C \emptyset + d_{k-1} P^n, & k = n+1 \end{cases}$$

So we have $[d_{k+1}, C] = d_k + \frac{1}{k!} \frac{\partial^k}{\partial \alpha^k} \Big|_{\alpha=0} \xi_\alpha$, $k \geq 0$. This can be reformulated in terms of $\Phi(t)$.

Proposition 11.1. *We have*

$$\Phi(t)C = (C + t)\Phi(t) + t\xi_t$$

Proof. Let P be an n -polytope. Then

$$\Phi(t)CP = CP + (CdP + P)t + \dots + (Cd_n P + d_{n-1} P)t^n + (1 + d_n P)t^{n+1} = C\Phi(t)P + t\Phi(t)P + t\xi_t P$$

□

Now let us consider the bipyramid. $dBP = 2CdP$ for any polytope of positive dimension, while $dB \text{ pt} = dI = 2 \text{ pt}$, and $2Cd \text{ pt} = 0$. So $dBP = 2CdP + 2\xi_0$. For $k \geq 2$ we have

$$d_k B P^n = \begin{cases} 2C d_k P^n + d_{k-1} P^n, & k < n+1; \\ 2 + d_{k-1} P^n = 2C \emptyset + d_{k-1} P^n, & k = n+1; \end{cases}$$

So $d_{k+1}B = 2Cd_{k+1} + d_k + \frac{2}{k!} \frac{\partial^k}{\partial \alpha^k} \Big|_{\alpha=0} \xi_\alpha$, $k \geq 1$.

Proposition 11.2.

$$\Phi(t)B = (B - 2C - t) + (2C + t)\Phi(t) + 2t\xi_t$$

Proof. For a point we have: $\Phi(t)B \text{ pt} = \Phi(t)I = I + (2 \text{ pt})t$, and

$$((B - 2C - t) + (2C + t)\Phi(t) + 2t\xi_t) \text{ pt} = I - 2I - (\text{pt})t + 2I + (\text{pt})t + 2(\text{pt})t = I + (2 \text{ pt})t$$

Let P be an n -polytope, $n > 0$. Then

$$\begin{aligned} \Phi(t)BP &= BP + (2CdP)t + (2Cd_2P + dP)t^2 + \cdots + (2Cd_nP + d_{n-1}P)t^n + (2 + d_nP)t^{n+1} = \\ &= (B - 2C - t)P + 2C\Phi(t)P + t\Phi(t)P + 2t\xi_tP \end{aligned}$$

□

Let us denote $A = 2C - B$. Then from Propositions 11.1 and 11.2 we have

$$\Phi(t)A = A + t + t\Phi(t)$$

At last, Propositions 11.1 and 11.2 imply the following formula.

Proposition 11.3.

$$\Phi(t)[B, C] = [B, C] + At + t^2$$

Proof.

$$\begin{aligned} \Phi(t)[B, C] &= \Phi(t)(BC - CB) = (B - 2C - t + (2C + t)\Phi(t) + 2t\xi_t)C - ((C + t)\Phi(t) + t\xi_t)B = \\ &= (B - 2C - t)C + (2C + t)((C + t)\Phi(t) + t\xi_t) + 2t^2\xi_t - (C + t)(B - 2C - t + (2C + t)\Phi(t) + 2t\xi_t) - t^2\xi_t = \\ &= (BC - CB) - t(B - 2C - t) + 2Ct\xi_t + t^2\xi_t + 2t^2\xi_t - 2Ct\xi_t - 2t^2\xi_t - t^2\xi_t = [B, C] + At + t^2 \end{aligned}$$

□

Now let us apply this formulas to the generalized f -polynomial.

Proposition 11.4. *Mappings $C, A: \mathcal{Qsym}[\alpha] \rightarrow \mathcal{Qsym}[\alpha]$ defined by the formulas*

$$Cg = (\alpha + \sigma_1)g + \sum_{n=1}^{\infty} t_n g(t_n, t_1, \dots, t_{n-1}, 0, 0, \dots)$$

$$Ag = \alpha g(\alpha, 0, 0, \dots) + t_1 g(\alpha, t_1, t_2, \dots) + \sum_{n=2}^{\infty} (t_n + t_{n-1})g(\alpha, t_n, t_{n+1}, \dots)$$

satisfy the relations

$$f(CP) = Cf(P), \quad f(AP) = Af(P)$$

for all $P \in \mathcal{P}$.

Proof. At first let us prove that this mappings are correctly defined, that is for any $g \in \mathcal{Qsym}[\alpha]$ its images Cg , and Ag belong to $\mathcal{Qsym}[\alpha]$. We will use the criterion obtained in Proposition 3.3 and Remark 3.4. It is easy to see that if g has bounded degree, then Cg and Ag have bounded degree.

Let $i \geq 1$. Then

$$\begin{aligned} (Cg)(\alpha, t_1, \dots, t_{i-1}, 0, t_{i+1}, \dots) &= (\alpha + \sum_{j \neq i} t_j)g(\alpha, t_1, \dots, t_{i-1}, 0, t_{i+1}, \dots) + \sum_{n=1}^{i-1} t_n g(t_n, t_1, \dots, t_{n-1}, 0, 0, \dots) + \\ &+ \sum_{n=i+1}^{\infty} t_n g(t_n, t_1, \dots, t_{i-1}, 0, t_{i+1}, \dots, t_{n-1}, 0, 0, \dots) = \\ &= (\alpha + \sigma_1(t_1, \dots, t_{i-1}, t_{i+1}, \dots))g(\alpha, t_1, \dots, t_{i-1}, t_{i+1}, \dots) + \sum_{n=1}^{i-1} t_n g(t_n, t_1, \dots, t_{n-1}, 0, 0, \dots) + \\ &+ t_{i+1}g(t_{i+1}, t_1, \dots, t_{i-1}, 0, 0, \dots) + \sum_{n=i+2}^{\infty} t_n g(t_n, t_1, \dots, t_{i-1}, t_{i+1}, \dots, t_{n-1}, 0, 0, \dots) = (Cg)(\alpha, t_1, \dots, t_{i-1}, t_{i+1}, \dots) \end{aligned}$$

Now consider A . For $i = 1$ the relation is clear. Let $i > 1$. Then

$$\begin{aligned}
(Ag)(\alpha, t_1, \dots, t_{i-1}, 0, t_{i+1}, \dots) &= \alpha g(\alpha, 0, 0, \dots) + \sum_{n=1}^{i-1} (t_n + t_{n-1}) g(\alpha, t_n, \dots, t_{i-1}, 0, t_{i+1}, \dots) + \\
&+ (0 + t_{i-1}) g(\alpha, 0, t_{i+1}, \dots) + (t_{i+1} + 0) g(\alpha, t_{i+1}, t_{i+2}, \dots) + \sum_{n=i+2}^{\infty} (t_n + t_{n-1}) g(\alpha, t_n, t_{n+1}, \dots) = \\
&= \alpha g(\alpha, 0, 0, \dots) + \sum_{n=1}^{i-1} (t_n + t_{n-1}) g(\alpha, t_n, \dots, t_{i-1}, t_{i+1}, \dots) + (t_{i+1} + t_{i-1}) g(\alpha, t_{i+1}, t_{i+2}, \dots) + \\
&+ \sum_{n=i+2}^{\infty} (t_n + t_{n-1}) g(\alpha, t_n, t_{n+1}, \dots) = (Ag)(\alpha, t_1, \dots, t_{i-1}, t_{i+1}, \dots)
\end{aligned}$$

On the other hand, we have

$$\begin{aligned}
\Phi(t_1)C &= (C + t_1)\Phi(t_1) + t_1\xi_{t_1} \\
\Phi(t_2)\Phi(t_1)C &= (C + t_2 + t_1)\Phi(t_2)\Phi(t_1) + t_2\xi_{t_2}\Phi(t_1) + t_1\xi_{t_1} \\
&\dots \\
\Phi(t_n)\Phi(t_{n-1})\dots\Phi(t_1)C &= (C + t_n + \dots + t_1)\Phi(t_n)\Phi(t_{n-1})\dots\Phi(t_1) + t_n\xi_{t_n}\Phi(t_{n-1})\dots\Phi(t_1) + \dots + t_1\xi_{t_1} \\
&\dots \\
\Phi_{\infty}C &= (C + \sigma_1)\Phi_{\infty} + \sum_{n=1}^{\infty} t_n f(t_n, t_1, \dots, t_{n-1}, 0, \dots) \\
f(CP) &= \xi_{\alpha}\Phi_{\infty}CP = (\alpha + \sigma_1)\xi_{\alpha}\Phi_{\infty}P + \sum_{n=1}^{\infty} t_n f(t_n, t_1, \dots, t_{n-1}, 0, \dots) = Cf(P) \\
&\dots \\
\Phi(t_1)A &= A + t_1 + t_1\Phi(t_1) \\
\Phi(t_2)\Phi(t_1)A &= A + t_2 + (t_2 + t_1)\Phi(t_2) + t_1\Phi(t_2)\Phi(t_1) \\
&\dots \\
\Phi(t_n)\Phi(t_{n-1})\dots\Phi(t_1)A &= A + t_n + (t_n + t_{n-1})\Phi(t_n) + (t_{n-1} + t_{n-2})\Phi(t_n)\Phi(t_{n-1}) + \dots + t_1\Phi(t_n)\dots\Phi(t_1) \\
&\dots \\
\Phi_{\infty}A &= A + t_1\Phi_{\infty}(t_1, t_2, \dots) + \sum_{n=2}^{\infty} (t_n + t_{n-1})\Phi_{\infty}(t_n, t_{n+1}, \dots) \\
f(AP) &= \xi_{\alpha}\Phi_{\infty}AP = \alpha f(\alpha, 0, 0, \dots) + t_1 f(\alpha, t_1, t_2, \dots) + \sum_{n=2}^{\infty} (t_n + t_{n-1}) f(\alpha, t_n, t_{n+1}, \dots) = Af(P)
\end{aligned}$$

□

Now the mapping $B : \mathcal{Qsym}[\alpha] \rightarrow \mathcal{Qsym}[\alpha]$ can be defined as $B = 2C - A$. Then $f(BP) = Bf(P)$.

11.2 The case of $\mathcal{RP}$

Now let us consider the ring $\mathcal{RP}$. In this ring CP is just the multiplication by the element $x = \text{pt}$. So

$$\begin{aligned}
\Phi(t)CP &= \Phi(t)(xP) = (\Phi(t)x) * (\Phi(t)P) = (x + t)\Phi(t)P = (C + t)\Phi(t)P \\
\Phi_{\infty}(CP) &= \Phi_{\infty}(xP) = \Phi_{\infty}(x) * \Phi_{\infty}(P) = (x + \sigma_1)\Phi_{\infty}(P) = (C + \sigma_1)\Phi_{\infty}(P) \\
f_{\mathcal{RP}}(CP) &= f_{\mathcal{RP}}(xP) = f_{\mathcal{RP}}(x)f_{\mathcal{RP}}(P) = (\alpha + \sigma_1)f_{\mathcal{RP}}(P) \\
F^*(CP) &= F^*(xP) = F^*(x)F^*(P) = \sigma_1 F^*(P)
\end{aligned}$$

In fact, the existence of the empty polytope $\emptyset$ such that $d_{n+1}P^n = \emptyset$ for any n -polytope P^n makes the formulas connecting C, B , and d_k simpler. We have:

$$\begin{aligned} [C, d] &= 1, \quad [C, d_{k+1}] = d_k, \quad k \geq 1 \\ dB &= 2Cd + \varepsilon_0, \quad d_{k+1}B = 2Cd_{k+1} + d_k, \quad k \geq 1 \end{aligned}$$

Proposition 11.5.

$$\Phi(t)B = (B - 2C - t) + (2C + t)\Phi(t) + \varepsilon_0 t$$

Proof. Indeed,

$$\begin{aligned} \Phi(t)BP &= BP + (2CdP + \varepsilon_0(P))t + (2Cd_2P + dP)t^2 + \sum_{k \geq 3} (2Cd_kP + d_{k-1}P)t^k = \\ &= (B - 2C - t)P + 2C\Phi(t)P + t\Phi(t)P + \varepsilon_0(P)t \end{aligned}$$

□

Then

$$\Phi(t)A = A + t + t\Phi(t) - \varepsilon_0 t$$

Proposition 11.6. *The mappings $A_{\mathcal{RP}}: \mathcal{Qsym}[\alpha] \rightarrow \mathcal{Qsym}[\alpha]$, $A_0: \mathcal{Qsym} \rightarrow \mathcal{Qsym}$ defined by the formulas*

$$\begin{aligned} A_{\mathcal{RP}}g &= \alpha g(\alpha, 0, 0, \dots) + t_1 g(\alpha, t_1, t_2, \dots) + \sum_{n=2}^{\infty} (t_n + t_{n-1})g(\alpha, t_n, t_{n+1}, \dots) - \sigma_1 g(0, 0, \dots) \\ A_0 g &= t_1 g(t_1, t_2, \dots) + \sum_{n=2}^{\infty} (t_n + t_{n-1})g(t_n, t_{n+1}, \dots) - \sigma_1 g(0, 0, \dots) \end{aligned}$$

satisfy the relations

$$f_{\mathcal{RP}}(AP) = A_{\mathcal{RP}}f_{\mathcal{RP}}(P), \quad F^*(AP) = A_0 F^*(P)$$

for all $P \in \mathcal{RP}$.

Proof. From the proof of Proposition 11.4 we see that both mappings $A_{\mathcal{RP}}$ and A_0 are correctly defined. Then

$$\begin{aligned} \Phi(t_1)A &= A + t_1 + t_1\Phi(t_1) - \varepsilon_0 t_1 \\ \Phi(t_2)\Phi(t_1)A &= A + t_2 + (t_2 + t_1)\Phi(t_2) + t_1\Phi(t_2)\Phi(t_1) - \varepsilon_0(t_2 + t_1) \\ &\dots \\ \Phi(t_n)\Phi(t_{n-1})\dots\Phi(t_1)A &= A + t_n + (t_n + t_{n-1})\Phi(t_n) + (t_{n-1} + t_{n-2})\Phi(t_n)\Phi(t_{n-1}) + \dots \\ &\quad + t_1\Phi(t_n)\dots\Phi(t_1) - \varepsilon_0(t_n + \dots + t_1) \\ &\dots \\ \Phi_{\infty}A &= A + t_1\Phi_{\infty}(t_1, t_2, \dots) + \sum_{n=2}^{\infty} (t_n + t_{n-1})\Phi_{\infty}(t_n, t_{n+1}, \dots) - \varepsilon_0\sigma_1 \\ f_{\mathcal{RP}}(AP) &= \varepsilon_{\alpha}\Phi_{\infty}A = \varepsilon_{\alpha}(AP) + t_1\varepsilon_{\alpha}\Phi_{\infty}(t_1, t_2, \dots) + \sum_{n=2}^{\infty} (t_n + t_{n-1})\varepsilon_{\alpha}\Phi_{\infty}(t_n, t_{n+1}, \dots) - \varepsilon_0(P)\sigma_1 = \\ &= \alpha f_{\mathcal{RP}}(\alpha, 0, 0, \dots) + t_1 f_{\mathcal{RP}}(\alpha, t_1, t_2, \dots) + \sum_{n=2}^{\infty} (t_n + t_{n-1})f_{\mathcal{RP}}(\alpha, t_n, t_{n+1}, \dots) - \sigma_1 f_{\mathcal{RP}}(0, 0, \dots) = A_{\mathcal{RP}}f_{\mathcal{RP}}(P) \\ F^*(AP) &= \varepsilon_0\Phi_{\infty}A = \varepsilon_0(AP) + t_1\varepsilon_0\Phi_{\infty}(t_1, t_2, \dots) + \sum_{n=2}^{\infty} (t_n + t_{n-1})\varepsilon_0\Phi_{\infty}(t_n, t_{n+1}, \dots) - \varepsilon_0(P)\sigma_1 = \\ &= t_1 F^*(t_1, t_2, \dots) + \sum_{n=2}^{\infty} (t_n + t_{n-1})F^*(t_n, t_{n+1}, \dots) - \sigma_1 F^*(0, 0, \dots) = A_0 F^*(P) \end{aligned}$$

□

As before $B_{\mathcal{RP}}$ and B_0 are defined as $B_{\mathcal{RP}} = 2C_{\mathcal{RP}} - A_{\mathcal{RP}}$, $B_0 = 2C_0 - A_0$, where $C_{\mathcal{RP}}g = (\alpha + \sigma_1)g$, $C_0g = \sigma_1g$, and

$$f_{\mathcal{RP}}(BP) = B_{\mathcal{RP}}f_{\mathcal{RP}}(P), \quad F^*(BP) = B_0F^*(B)$$

At last

$$\begin{aligned} \Phi(t)(BC - CB) &= (B - 2C - t + (2C + t)\Phi(t) + \varepsilon_0t)C - (C + t)\Phi(t)B = \\ &= BC - (2C + t)C + (2C + t)(C + t)\Phi(t) - (C + t)(B - 2C - t + (2C + t)\Phi(t) + \varepsilon_0t) = \\ &= BC - CB + (2C - B)t + t^2 - (C + t)\varepsilon_0t = BC - CB + At + t^2 - (x + t)t\varepsilon_0 \end{aligned}$$

So we obtain the following proposition.

Proposition 11.7. *In the ring $\mathcal{RP}$*

$$\Phi(t)[B, C] = [B, C] + At + t^2 - (x + t)t\varepsilon_0$$

12 Problem of the Description of Flag Vectors of Polytopes

We have mentioned that on the space of simple polytopes

$$f(\alpha, t_1, t_2, \dots)(P^n) = f_1(\alpha, t_1 + t_2 + \dots)(P^n) = f_1(\alpha, \sigma_1)(P^n).$$

The only linear relation on the polynomial f_1 is $f_1(-\alpha, \alpha + t) = f_1(\alpha, t)$, and it is equivalent to the Dehn-Sommerville relations. In fact, for the polynomial $g = g(\alpha, \sigma_1) \in \text{Qsym}[t_1, t_2, \dots][\alpha]$ this condition is necessary and sufficient to be an image of an integer combination of simple polytopes.

One of the outstanding results in the polytope theory is the so-called g -theorem, which was formulated as a conjecture by P. McMullen [Mc1] in 1970 and proved by R. Stanley [St1] (the necessity) and L. Billera and C. Lee [BL] (the sufficiency) in 1980.

For an n -dimensional simple polytope P^n let us define an h -polynomial:

$$h(\alpha, t)(P^n) = h_0\alpha^n + h_1\alpha^{n-1}t + \dots + h_{n-1}\alpha t^{n-1} + h_n t^n = f_1(\alpha - t, t)(P^n)$$

Since $f_1(-\alpha, \alpha + t) = f_1(\alpha, t)$, the h -polynomial is symmetric: $h(\alpha, t) = h(t, \alpha)$. So $h_i = h_{n-i}$.

Definition 12.1. A g -vector of a simple polytope P^n is the set of numbers $g_0 = 1$, $g_i = h_i - h_{i-1}$, $1 \leq i \leq [\frac{n}{2}]$.

For any positive integers a and i there exists a unique "binomial i -decomposition" of a :

$$a = \binom{a_i}{i} + \binom{a_{i-1}}{i-1} + \dots + \binom{a_j}{j},$$

where $a_i > a_{i-1} > \dots > a_j \geq j \geq 1$.

Let us define

$$a^{(i)} = \binom{a_i + 1}{i + 1} + \binom{a_{i-1} + 1}{i} + \dots + \binom{a_j + 1}{j + 1}, \quad 0^{(i)} = 0.$$

For example, $a = \binom{a}{1}$, so $a^{(1)} = \binom{a+1}{2} = \frac{(a+1)a}{2}$; for $i \geq a$ we have: $a = \binom{i}{i} + \binom{i-1}{i-1} + \dots + \binom{i-a+1}{i-a+1}$, so $a^{(i)} = a$.

Theorem (g -theorem, [St1], [BL]). *Integer numbers $(g_0, g_1, \dots, g_{[\frac{n}{2}]})$ form the g -vector of some simple n -dimensional polytope if and only if they satisfy the following conditions:*

$$g_0 = 1, \quad 0 \leq g_1, \quad 0 \leq g_{i+1} \leq g_i^{(i)}, \quad i = 1, 2, \dots, \left[\frac{n}{2}\right] - 1.$$

As it is mentioned in [Z1] even for 4-dimensional non-simple polytopes the corresponding problem involving flag f -vectors is extremely hard.

Modern tools used to obtain linear and nonlinear inequalities satisfied by flag f -numbers are based on the notion of a cd -index [BK, BE1, BE2, EF, ER, L, St3], a *toric h -vector* [BE, BL, K, St2, Sts] and its generalizations [F1, F2, L], a *ring of chain operators* [BH, BLiu, Kalai].

In [St1] R. Stanley constructed for a simple polytope P a projective toric variety X_P so that $\{h_i\}$ are the even Betti numbers of the (singular) cohomology of X_P . Then the Poincare duality and the Hard Lefschetz theorem for X_P imply the McMullen conditions for P .

In [Mc2] P. McMullen gave the purely geometric proof of these conditions using the notion of a *polytope algebra*.

In [St2] Stanley generalized the definition of an h -vector to an arbitrary polytope P in such a way that if a polytope is rational the generalized $\hat{h}_i$ are the intersection cohomology Betti numbers of the associated toric variety (see the validation of the claim in [Fi]).

The set of numbers $(\hat{h}_0, \dots, \hat{h}_n)$ is called a *toric h -vector*. It is nonnegative and symmetric: $\hat{h}_i = \hat{h}_{n-i}$, and in the case of rational polytope the Hard Lefschetz theorem in the intersection cohomology of the associated toric variety proves the unimodality

$$1 = \hat{h}_0 \leq \hat{h}_1 \leq \dots \leq \hat{h}_{\lfloor \frac{n}{2} \rfloor}$$

Generalizing the geometric method introduced by P. McMullen [Mc2] Kalle Karu [K] proved the unimodality for an arbitrary polytope.

The toric h -vector consists of linear combinations of the flag f -numbers. In general case it does not contain the full information about the flag f -vector. In [L] C. Lee introduced an "extended toric" h -vector that carries all the information about the flag f -numbers and consists of a collection of nonnegative symmetric unimodal vectors. In [F1] and [F2] J. Fine introduced the generalized h - and g -vectors that contain the toric h - and g -vectors as subsets, but can have negative entries.

Let us look on the problem of the description of flag f -vectors from the point of view of the ring of polytopes.

g -theorem describes all the polynomials $\psi(\alpha, t_1, t_2, \dots) \in \text{Qsym}[t_1, t_2, \dots][\alpha]$ that are images of simple polytopes under the ring homomorphism f :

$$\psi(\alpha, t_1, t_2, \dots) = f(\alpha, t_1, t_2, \dots)(P^n), \quad P^n - \text{a simple } n\text{-dimensional polytope.}$$

In the general case we know the criterion when the polynomial $\psi \in \text{Qsym}[t_1, t_2, \dots][\alpha]$ is the image of an integer combination of polytopes.

Question 12.2. Find a criterion when the polynomial $\psi \in \text{Qsym}[t_1, t_2, \dots][\alpha]$ is the image of an n -dimensional polytope.

A Join

Proposition A.1. $P^{n_1} * Q^{n_2}$ is an $(n_1 + n_2 + 1)$ -dimensional polytope.

Faces of $P^{n_1} * Q^{n_2}$ up to an affine equivalence are exactly $F * G$, where $\emptyset \subseteq F \subseteq P^{n_1}$, $\emptyset \subseteq G \subseteq Q^{n_2}$ are faces of P^{n_1} and Q^{n_2} respectively.

Proof. The formula for dimension of $P^{n_1} * Q^{n_2}$ follows from Remark 2.13.

The nonempty faces of a polytope are exactly the subsets of the polytope that minimize some linear function on it. Let $\mathbf{c} = (c_1, c_2)$ be a linear function in $(\mathbb{R}^{n_1+1} \times \mathbb{R}^{n_2+1})^* = (\mathbb{R}^{n_1+1})^* \times (\mathbb{R}^{n_2+1})^*$. Then c_1 has its minimum c_1 on the face F of P^{n_1} , and c_2 has its minimum c_2 on the face G of Q^{n_2} . Since any point of $P^{n_1} * Q^{n_2}$ has the form $t_1 \mathbf{p} + t_2 \mathbf{q}$, where $\mathbf{p} \in P^{n_1}$, $\mathbf{q} \in Q^{n_2}$, $t_1, t_2 \geq 0$, $t_1 + t_2 = 1$, we have

- if $c_1 < c_2$, then $\mathbf{c}$ has its minimum on the face $F * \emptyset$ of $P^{n_1} * Q^{n_2}$;
- if $c_1 > c_2$, then $\mathbf{c}$ has its minimum on $\emptyset * G$;
- if $c_1 = c_2$, then $\mathbf{c}$ has its minimum on the set

$$X = \{t_1 \mathbf{p} + t_2 \mathbf{q}, \mathbf{p} \in F, \mathbf{q} \in G, t_1, t_2 \geq 0, t_1 + t_2 = 1\}.$$

Let F lie in the k -simplex $\text{conv}\{\mathbf{p}_1, \dots, \mathbf{p}_{k+1}\} \subset \text{aff}(F)$, and G lie in the l -simplex $\text{conv}\{\mathbf{q}_1, \dots, \mathbf{q}_{l+1}\} \subset \text{aff}(G)$. Now let $\{\overrightarrow{O\mathbf{p}_i}, i = 1, \dots, k+1\}$ be a basis in $\mathbb{R}^{k+1} = \text{aff}\{O, \mathbf{p}_i, i = 1, \dots, k+1\}$ and $\{\overrightarrow{O\mathbf{q}_j}, j = 1, \dots, l+1\}$ be a basis in $\mathbb{R}^{l+1} = \text{aff}\{O, \mathbf{q}_j, j = 1, \dots, l+1\}$. Then $X = F * G$.

Note that $\mathbf{c}_1$ can have the constant value on P^{n_1} , or $\mathbf{c}_2$ can have the constant value on Q^{n_2} . In these cases $F = P^{n_1}$, or $G = Q^{n_2}$.

Let λ_P be the linear function equal to the sum of all the coordinates in $\mathbb{R}^{n_1+1}$, and λ_Q be the linear function equal to the sum of all the coordinates in $\mathbb{R}^{n_2+1}$. Then

$$\begin{aligned} \lambda_P(\mathbf{p}) &= 1, & \lambda_Q(\mathbf{p}) &= 0 & \text{for any } \mathbf{p} \in P^{n_1} = P^{n_1} * \emptyset \\ \lambda_P(\mathbf{q}) &= 0, & \lambda_Q(\mathbf{q}) &= 1 & \text{for any } \mathbf{q} \in Q^{n_2} = \emptyset * Q^{n_2} \end{aligned}$$

Now let $F \neq \emptyset$, $F \subseteq P^{n_1}$ be a face of P^{n_1} , and let $\mathbf{c}_1 \in (\mathbb{R}^{n_1+1})^*$ be a linear function that has its minimum c_1 exactly on F . Similarly, let $(G, \mathbf{c}_2, c_2)$ be the corresponding data for Q^{n_2} . Then let $\mathbf{c} = (\mathbf{c}_1, \mathbf{c}_2) \in (\mathbb{R}^{n_1+1} \times \mathbb{R}^{n_2+1})^*$. By an addition or a subtraction of λ_P and λ_Q with coefficients we can obtain any of the cases $c_1 < c_2$, $c_1 = c_2$, or $c_1 > c_2$, so $F * \emptyset$, $F * G$, $\emptyset * G$ are faces of $P^{n_1} * Q^{n_2}$.

Thus faces of the polytope $P^{n_1} * Q^{n_2}$ up to an affine equivalence have the form

$$F * G, \emptyset \subseteq F \subseteq P^{n_1}, \emptyset \subseteq G \subseteq Q^{n_2}.$$

□

B Structure theorems

Proposition B.1. *$\mathcal{P}$ is a polynomial ring generated by indecomposable combinatorial polytopes.*

Proof. We will show that any polytope P^n of positive dimension can be represented in a unique up to the order of factors way as a product of indecomposable polytopes of positive dimensions $P = P_1 \times \cdots \times P_k$.

It is clear that such a decomposition exists. Now let

$$P_1 \times \cdots \times P_k = Q_1 \times \cdots \times Q_s,$$

where

$$1 \leq \dim P_1 \leq \dim P_2 \leq \cdots \leq \dim P_k, \quad \text{and} \quad 1 \leq \dim Q_1 \leq \dim Q_2 \leq \cdots \leq \dim Q_s.$$

Let us consider a face F of the form $\mathbf{v}_1 \times \cdots \times \mathbf{v}_{k-1} \times P_k \simeq P_k$, where $\mathbf{v}_i$ are vertices. Then we have $F = G_1 \times \cdots \times G_s$, where G_i is a face of Q_i . Since P_k is indecomposable, we see that all G_j but one are points, and P_k is combinatorially equivalent to a face of some Q_r . Similarly Q_r is combinatorially equivalent to a face of some P_t . Then $\dim P_k \leq \dim Q_r \leq \dim P_t \leq \dim P_k$, therefore, $P_k = Q_r$. Similarly $Q_s \simeq P_l$ for some l . So $\dim P_k = \dim Q_s$ and without loss of generality we can set $P_k = Q_s$.

If $P_k = I$ then $P_i = Q_j = I$ for all i, j , so the statement is true. Let $P_k \neq I$. We know that

$$\mathbf{v}_1 \times \cdots \times \mathbf{v}_{k-1} \times P_k = \mathbf{w}_1 \times \cdots \times \mathbf{w}_{s-1} \times P_k$$

Let us consider a face G of the form $\mathbf{v}_1 \times \cdots \times \mathbf{v}_{i-1} \times I \times \mathbf{v}_{i+1} \times \cdots \times \mathbf{v}_{k-1} \times P_k$, where I is the edge that joins the vertices $\mathbf{v}_i$ and $\mathbf{v}'_i$ of the polytope P_i . Then $G = \mathbf{w}_1 \times \cdots \times \mathbf{w}_{j-1} \times J \times \mathbf{w}_{j+1} \times \cdots \times \mathbf{w}_{s-1} \times P_k$, where J is the edge that joins the vertices $\mathbf{w}_j$ and $\mathbf{w}'_j$ of some polytope Q_j .

Since $\mathbf{v}_1 \times \cdots \times \mathbf{v}'_i \times \cdots \times P_k \subset G$, P_k is indecomposable, and $P_k \neq J$, we have

$$\mathbf{v}_1 \times \cdots \times \mathbf{v}'_i \times \cdots \times P_k = \mathbf{w}_1 \times \cdots \times \mathbf{w}'_j \times \cdots \times P_k.$$

Changing the edges we can achieve any vertex $\mathbf{v}'_1 \times \cdots \times \mathbf{v}'_{k-1}$ of $P_1 \times \cdots \times P_{k-1}$, so for any such a vertex we have $\mathbf{v}'_1 \times \cdots \times \mathbf{v}'_{k-1} \times P_k = \mathbf{w}'_1 \times \cdots \times \mathbf{w}'_{s-1} \times P_k$. Therefore for any face $F_1 \times \cdots \times F_{k-1} \times P_k$

$$F_1 \times \cdots \times F_{k-1} \times P_k = G_1 \times \cdots \times G_{s-1} \times P_k$$

Thus for any face F of the polytope $P_1 \times \cdots \times P_{k-1}$ there is a face G of the polytope $Q_1 \times \cdots \times Q_{s-1}$ such that $F \times P_k = G \times P_k$, and vice versa. Moreover, $F \subset F'$ if and only if $G \subset G'$. Thus

$$P_1 \times \cdots \times P_{k-1} = Q_1 \times \cdots \times Q_{s-1}$$

Iterating this step in the end we obtain that $k = s$, and up to the order of the polytopes $P_i = Q_i$, $i = 1, \dots, k$. □

Proposition B.2. $\mathcal{RP}$ is a ring of polynomials generated by all join-indecomposable polytopes.

Proof. Let $P_1 * P_2 \cdots * P_k = Q_1 * Q_2 * \cdots * Q_s$, where P_i, Q_j are polytopes,

$$0 \leq \dim P_1 \leq \dim P_2 \leq \cdots \leq \dim P_k, \quad 0 \leq \dim Q_1 \leq \dim Q_2 \leq \cdots \leq \dim Q_s$$

and each polytope P_i and Q_j can not be decomposed as a join of polytopes of lower dimensions. Let $\dim P_k \geq \dim Q_s$. Consider the face $\emptyset * \emptyset * \cdots * P_k$. It can be represented in the form $G_1 * G_2 * \cdots * G_s$, where G_j is a face of Q_j . Since P_k is join-indecomposable, all G_j but one should be equal to $\emptyset$, and one of them should be equal to P_k . Let $P_k = G_r$. Then P_k is a face of Q_r , so $P_k = Q_r$, since $\dim P_k \geq \dim Q_s \geq \dim Q_r$. Without loss of generality we can take $r = s$.

We have $P' * P_k = Q' * P_k$, and the combinatorial equivalence identifies the face $\emptyset * P_k$ of the first polytope with the same face $\emptyset * P_k$ of the second. Consider the face $P' * \emptyset$. It should have the form $G_1 * G_2$ where G_1 is a face of Q' and G_2 is a face of P_k . Then $\emptyset * G_2$ should be a face of $\emptyset * P_k$ and $P' * \emptyset$, therefore $G_2 = \emptyset$ and $G_1 = P'$. Since $\dim P' = \dim Q'$, we see that $P' = Q'$.

So $P_1 * P_2 * \cdots * P_{k-1} = Q_1 * Q_2 * \cdots * Q_{s-1}$. Let $s \geq k$. Iterating this step in the end we obtain $P_1 = Q_1 * Q_2 * \cdots * Q_{s-k+1}$. Therefore $k = s$ and $P_1 = Q_1$.

Thus each polytope can be decomposed in a unique up to permutations way as $P_1 * \cdots * P_k$, where P_j are join-indecomposable. Since $\mathcal{RP}$ additively is a free abelian group generated by combinatorial polytopes and $\emptyset$, it is a ring of polynomials generated by all join-indecomposable polytopes. $\square$

C Examples

Proposition C.1. Any monomial M_ω such that $M_\omega \in F(\mathcal{RP})$ has the form $M_{(2k+1)}$, $k \geq 0$.

Proof. Indeed, for the monomial $M_{(a_1, \dots, a_k)}$ we have

$$M_{(a_1, \dots, a_k)}(t, -t, t_1, t_2, \dots) = M_{(a_1, \dots, a_k)}(t_1, t_2, \dots) + (t^{a_1} + (-t)^{a_1}) M_{(a_2, \dots, a_k)}(t_1, t_2, \dots) + (-1)^{a_2} t^{a_1+a_2} M_{(a_3, \dots, a_k)}$$

If $k \geq 2$ the last summand does not cancel, so $k = 1$. For $M_{(n)}$ with $n = 2l + 1$ we have

$$M_{(2l+1)}(t_1, \dots, t_{i-1}, t, -t, t_i, \dots) = t_1^{2l+1} + \cdots + t_{i-1}^{2l+1} + t^{2l+1} - t^{2l+1} + t_i^{2l+1} + \cdots = M_{(2l+1)}(t_1, \dots, t_{i-1}, t_i, \dots),$$

while for $n = 2l$ the term containing $t^{2l} + (-t)^{2l}$ does not cancel. $\square$

Proposition C.2. $M_{(2k+1, 2l+1)} - M_{(2l+1, 2k+1)} \in F(\mathcal{RP})$

Proof. Indeed,

$$M_{(2k+1, 2l+1)}(t_1, \dots, t_{i-1}, t, -t, t_i, \dots) = M_{(2k+1, 2l+1)}(t_1, \dots, t_{i-1}, t_i, \dots) + \sum_{j < i} t_j^{2k+1} (t^{2l+1} + (-t)^{2l+1}) + \sum_{j \geq i} (t^{2k+1} + (-t)^{2k+1}) t_j^{2l+1} + (-1)^{2l+1} t^{2k+2l+2}$$

Then

$$(M_{(2k+1, 2l+1)} - M_{(2l+1, 2k+1)})(t_1, \dots, t_{i-1}, t, -t, t_i, \dots) = (M_{(2k+1, 2l+1)} - M_{(2l+1, 2k+1)})(t_1, \dots, t_{i-1}, t_i, \dots)$$

$\square$

This proposition can be generalized.

Proposition C.3. Let $(j_1, \dots, j_k)$ be the set of odd positive integer numbers. Then the quasi-symmetric polynomial

$$\sum_{\sigma \in S_k} (-1)^\sigma M_{(j_{\sigma(1)}, \dots, j_{\sigma(k)})}$$

belongs to the image of F .

Proof. Indeed, since j_l are odd, we have

$$M_{(j_1, \dots, j_k)}(t_1, \dots, t_{i-1}, t, -t, t_i, \dots) = M_{(j_1, \dots, j_k)}(t_1, \dots, t_{i-1}, t_i, \dots) - \sum_{l_1 < \dots < l_s < i \leq l_{s+3} < \dots < l_k} t_{l_1}^{j_1} \dots t_{l_s}^{j_s} t^{j_{s+1} + j_{s+2}} t_{l_{s+3}}^{j_{s+3}} \dots t_{l_k}^{j_k}$$

For any pair (j_{s+1}, j_{s+2}) there exists a unique monomial $M_{(j_1, \dots, j_s, j_{s+2}, j_{s+1}, j_{s+3}, \dots, j_k)}$, and this monomial has the opposite sign. Therefore all the monomials of the form $t_{l_1}^{j_{\sigma(1)}} \dots t_{l_s}^{j_{\sigma(s)}} t^{j_{\sigma(s+1)} + j_{\sigma(s+2)}} t_{l_{s+3}}^{j_{\sigma(s+3)}} \dots t_{l_k}^{j_{\sigma(k)}}$ vanish. $\square$

Let us calculate the image of the Ehrenborg transformation F in small dimensions.

$n = 0$ The image of the point $x = \text{pt}$ is $M_{(1)} = \sigma_1$. So

$$F(\mathcal{RP}^2) = \mathcal{R}^*((\mathcal{D}^*)^2) = \mathbb{Z}\langle\sigma_1\rangle.$$

$n = 1$ The image of the segment $I = \text{pt} * \text{pt}$ is $M_{(2)} + 2M_{(1,1)} = \sigma_1^2$. So

$$F(\mathcal{RP}^4) = \mathcal{R}*((\mathcal{D}^*)^4) = \mathbb{Z}\langle\sigma_1^2\rangle.$$

$n = 2$ The image of the m -gon M_m^2 is

$$M_{(3)} + mM_{(2,1)} + mM_{(1,2)} + 2mM_{(1,1,1)} = M_{(3)} + m(\sigma_1\sigma_2 - \sigma_3)$$

Therefore,

$$F(\mathcal{RP}^6) = \mathcal{R}*((\mathcal{D}^*)^6) = \mathbb{Z}\langle M_{(3)}, \sigma_1\sigma_2 - \sigma_3 \rangle,$$

where

$$M_{(3)} = F(4\Delta^2 - 3I^2), \quad \sigma_1\sigma_2 - \sigma_3 = F(I^2 - \Delta^2),$$

$n = 3$ The image of a 3-dimensional polytope is

$$M_{(4)} + f_0M_{(1,3)} + f_1M_{(2,2)} + f_2M_{(3,1)} + f_{01}M_{(1,1,2)} + f_{02}M_{(1,2,1)} + f_{12}M_{(2,1,1)} + f_{012}M_{(1,1,1,1)}$$

Since $f_0 - f_1 + f_2 = 2$, $f_{01} = f_{02} = f_{12} = 2f_1$, and $f_{012} = 4f_1$, we obtain

$$(M_{(4)} + 2M_{(3,1)}) + f_0(M_{(1,3)} - M_{(3,1)}) + f_1(M_{(2,2)} + M_{(3,1)} + 2(\sigma_1\sigma_3 - 2\sigma_4)),$$

Then

$$F(\mathcal{RP}^8) = \mathcal{R}*((\mathcal{D}^*)^8) = \mathbb{Z}\langle M_{(4)} + 2M_{(3,1)}, M_{(1,3)} - M_{(3,1)}, M_{(2,2)} + M_{(3,1)} + 2(\sigma_1\sigma_3 - 2\sigma_4) \rangle,$$

where

$$M_{(4)} + 2M_{(3,1)} = F(5\Delta^3 - 6CI^2 + 2B\Delta^2), \quad M_{(1,3)} - M_{(3,1)} = F(-\Delta^3 + 3CI^2 - 2B\Delta^2) \\ M_{(2,2)} + M_{(3,1)} + 2(\sigma_1\sigma_3 - 2\sigma_4) = F(-CI^2 + B\Delta^2)$$

References

- [ABS] M. Aguiar, N. Bergeron, F. Sottile, *Combinatorial Hopf Algebras and Generalized Dehn-Sommerville relations*. Compositio Mathematica 142(1) 2006, 1–30, arXiv: math/0310016v1 [math.CO].
- [BR] A. Baker, B. Richter, *Quasisymmetric functions from a topological point of view*. Math. Scand. 103 (2008), 208–242.
- [BB] M. M. Bayer, L. J. Billera, *Generalized Dehn-Sommerville relations for polytopes, spheres and Eulerian partially ordered sets*. Invent. Math. 79: 143–157, 1985.

- [BE] M. M. Bayer and R. Ehrenborg, *The toric h -vectors of partially ordered sets*. Trans. Amer. Math. Soc. 352 (2000), 4515–4531 (electronic).
- [BK] M. M. Bayer, A. Klapper, *A new index for polytopes*. Discrete Comput. Geom. 6, 33–47 (1991).
- [BL] M. M. Bayer and C. W. Lee, *Combinatorial aspects of convex polytopes*. in Handbook of convex geometry, Vol. A, B, North-Holland, Amsterdam, 1993, pp. 485–534.
- [BMSW] N. Bergeron, S. Mykytiuk, F. Sottile, and S. van Willigenburg, *Pieri Operations on Posets*. J. of Comb. Theory Series A 91(2000) 84–110. arXiv: math/0002073v2 [math.CO] 21 Jul 2000.
- [BE1] L. Billera, R. Ehrenborg, *Monotonicity of the cd -index for polytopes*. Math. Z. 233(200) 421–441.
- [BE2] L. Billera, R. Ehrenborg and M. Readdy, *The cd -index of zonotopes and arrangements*. Mathematical Essays in Honor of Gian-Carlo Rota, Birkhauser, Boston, 1998.
- [BH] L. Billera, G. Hettley, *Linear inequalities for flags in graded posets*. J. of Combinatorial Theory Series A, 2000, V89, N1, 77–104, arXiv: math/9706220v1 [math.CO]
- [BL] L. Billera, C. Lee, *A proof of sufficiency of McMullen’s conditions for f -vectors of simplicial polytopes*. J. Combin. Theory, Ser. A, 1981, V31, N3, 237–255.
- [BLiu] L. Billera, N. Liu, *Non-commutative enumeration in graded posets*. Journal of Algebraic Combinatorics 12 (2000), 7–24.
- [BS] R. Bott, H. Samelson, *On the Pontryagin product in spaces of paths*. Comment. Math. Helv., 27, 1953, 320–337
- [BP] V. M. Buchstaber, T. E. Panov, *Torus actions and their applications in topology and combinatorics*, Providence, R.I.: American Mathematical Society, 2002. (University Lecture Series; V.24).
- [BR] V. M. Buchstaber, N. Ray, *An invitation to toric topology: Vertex four of a remarkable tetrahedron*. Toric topology. Contemporary Mathematics (AMS), v. 460, Providence, R.I., 2008, 1–27; MIMS, 2008.31: <http://www.manchester.ac.uk/mims/eprints>.
- [Buch] V. M. Buchstaber, *Ring of Simple Polytopes and Differential Equations*. Proceedings of the Steklov Institute of Mathematics, v. 263, 2008, 1–25.
- [BuchE] Victor M. Buchstaber, Nikolai Erokhovets, *Ring of Polytopes, Quasi-symmetric functions and Fibonacci numbers*, arXiv:1002.0810v1 [math.CO].
- [BuchE1] V. M. Buchstaber, N. Erokhovets *Algebra of operators on the ring of polytopes and quasi-symmetric functions*, Russian Mathematical Surveys (2010), 65(2):381.
- [C] H. S. M. Coxeter, *Regular Polytopes*. 3rd edition, Dover Publications, 1973.
- [CFL] K. T. Chen, R. H. Fox, and R. C. Lyndon, *Free differential calculus*. IV, Ann. Math. 68 (1958), 81–95.
- [Ehr] R. Ehrenborg, *On Posets and Hopf Algebras*. Advances in Mathematics, 119, 1–25 (1996).
- [EF] R. Ehrenborg, H. Fox, *Inequalities for cd -indices of joins and products of polytopes*. Combinatorica 23 (2003), 427–452.
- [EKZ] D. Eppstein, G. Kuperberg, G. M. Ziegler, *Fat 4-polytopes and fatter 3-spheres*. Discrete Geometry: in honour of W. Kuperberg’s 60th birthday, Pure and Applied Mathematics. A series of Monographs and Textbooks, vol. 253, Marcel Dekker, Inc., 2003, pp.239–265.
- [ER] R. Ehrenborg, M. Readdy, *Coproducts and the cd -index*. J. Algebraic Combin. 8 (1998), 273–299.
- [Fi] K. -H. Fieseler, *Rational Intersection Cohomology of Projective Toric Varieties*. Journal f.d. reine u. angew. Mathematik 413 (1991), 88–98.

- [F1] J. Fine, *A complete h-vector for convex polytopes*. Preprint, arXiv: 0911.5722v1 [math.CO]
- [F2] J. Fine, *A complete g-vector for convex polytopes*. Preprint, arXiv: 1001.1562v1 [math.CO]
- [Fr] A. Frohmader, *Face vectors of flag complexes*. Israel Journal of Mathematics, 164, 2008, 153–164.
- [Gb] B. Grunbaum, *Convex Polytopes*. Vol. 221 of Graduate Texts in Mathematics, Springer-Verlag, New York, Second ed., 2003.
- [H] M. Hazewinkel, *The Algebra of Quasi-Symmetric Functions is free over integers*. Advances in Mathematics 164, 283–300 (2001).
- [JR] S. Joni and G.-C. Rota, *Coalgebras and bialgebras in combinatorics*. Stud. Appl. Math. 61(1979), 93–139.
- [Kalai] G. Kalai, *A new basis for polytopes*. J. Combinatorial Theory, Ser A 49 (1988), 191–208.
- [K] Kalle Karu, *Hard Lefschetz Theorem for Nonrational Polytopes*. Invent. Math. 157 (2004), 419–447, arXiv: math/0112087v4 [math.AG] 17 Dec 2002.
- [L] Carl W. Lee, *Sweeping the cd-Index and the Toric h-Vector*. 2009, preprint: <http://www.ms.uky.edu/~lee/cd.pdf>
- [Mc1] P. McMullen, *The numbers of faces of simplicial polytopes*. Israel J. Math. 1971, V9, 559–570.
- [Mc2] P. McMullen, *On Simple Polytopes*. Invent. math. 113, 419–444 (1993).
- [N] S. P. Novikov, *Various doublings of Hopf algebras. Operator algebras on quantum groups, complex cobordisms*. Russian Mathematical Surveys, 47:5, 1992, 198–199.
- [RS] N. Ray, W. Schmitt, *Combinatorial models for coalgebraic structures*. Advances in Mathematics 1998, Vol 138, Iss 2, 211–262.
- [Re] C. Reutenauer, *Free Lie algebras*. Oxford Univ. Press, Oxford, UK, 1993.
- [Sch1] William Schmitt, *Antipodes and incidence coalgebras*. Journal of Combinatorial Theory A, 46:264–290, 1987.
- [Sch2] William Schmitt, *Incidence Hopf Algebras*. J. Pure Appl. Algebra 96(1994), 299–330.
- [St1] R. P. Stanley, *The number of faces of simplicial convex polytope*. Advances in Math. 1980. V. 35, N3, 236–238.
- [St2] R. P. Stanley, *Generalized h-vectors, intersection cohomology of toric varieties, and related results*. Commutative Algebra and Combinatorics, Advanced Studies in Pure Mathematics 11, Kinokuniya, Tokyo, and North-Holland, Amsterdam/New York, 1987, pp. 187–213.
- [St3] R. P. Stanley, *Flag f-vectors and the cd-index*. Math. Z. 216, 483–499 (1994).
- [Sts] C. Stenson, *Relationships among flag f-vector inequalities for polytopes*. Discrete and Computational Geometry 31 (2004), pp. 257–273
- [Z1] G. M. Ziegler, *Face numbers of 4-polytopes and 3-spheres*. Proceedings of the ICM 2002 Beijing, Volume III, 2004, 625–634; arXiv: math/0208073v2 [math.MG] 20 Jun 2003.
- [Z2] G. M. Ziegler, *Lectures on Polytopes*. Springer-Verlag, New York, 2007.